

AUGUST 1, 1950


# MODERN MEDICINE

*The Journal of Diagnosis and Treatment*



Dr. Robert L. Bennett  
*(see page 11)*

*Table of contents*  
*page 11*



Just a small dose provides

## **Round-the- clock relief for allergic patients**

Long-lasting relief<sup>1</sup> . . . low milligram dosage<sup>2</sup> with few side actions. These are advantages you *expect* in a professional antihistamine for use in difficult allergies, or for patients who have not responded to other drugs. These advantages are *combined* in Decapryn—the antihistamine that provides a good night's rest and a good day's work for allergic patients.

1 "Symptoms were relieved from 4 to 24 hours after the administration of a single dose of Decapryn—" . . . Sheldon, J.M. Et al; Univ. Mich. Hosp. Bull. 11:13-15 (1939)

2 "It was found that 12.5 mg. could be given during the day with comparatively few side reactions and yet maintain good clinical results—" . . . MacQuibdy, E.L.; Neb. State M.J. 31:123 (1939)

# DECAPRYN<sup>®</sup>

The long-lasting, low-dosage prescription antihistamine

DECAPRYN (DOXYLAMINE) SUCCINATE



Available on prescription only, as pleasant-tasting liquid, or palatable tablets



In obesity, where control of appetite is desired  
... in mild depression, drowsiness ... and  
to overcome sedative action of antihistaminics ...  
as adjunctive treatment in acute and chronic alcoholism

# SYNDROX\*

(Brand of Methamphetamine Hydrochloride)

- ....elevates the mood (produces euphoria)
- ....imparts a sense of increased energy and efficiency
- ....counteracts sleepiness and feeling of fatigue
- ....suppresses appetite



## ADVANTAGES OF SYNDROX:

- Rapid onset (10-20 minutes)
- Long duration of effect (6-12 hours, depending on dose)
- Negligible side effects, with proper dosage
- Small dosage

**Suggested initial dose:** 2.5 to 5 mg. daily; dosage may be increased to 2.5 to 5 mg. two to three times daily and maintained at this level as long as there are no untoward effects.

\*TRADEMARK OF MCNEIL LABORATORIES, INC.

*Supplied in 5 mg. tablets (scored, green), bottles of 100 and 1000. Also available in a pleasant-tasting elixir (colored amber); each 30 cc. (1 fl. oz.) containing 20 mg.—pints and gallons. Samples on request.*

# McNEIL

LABORATORIES, INC. PHILADELPHIA 32, PENNSYLVANIA



## NOW, BRIGHTER PROSPECTS FOR THE CARDIAC PATIENT

New hope for the cardiac patient's future comes from today's more efficient management. Theorate 'Robins' helps tide over the coronary patient, by its potent coronary vasodilation, nontoxic diuresis and effective myocardial stimulation. It may be used over long periods without fear of gastric irritation or oversedation. Theorate 'Robins' is recommended for coronary disease, angina pectoris, congestive heart failure, essential hypertension, edema and other cardiovascular pathologies. • Each enteric-coated tablet contains: 100% alkaloidal Theobromine 5 gr., Phenobarbital ¼ gr.

A. H. ROBINS COMPANY, INC. • Richmond 20, Va.  
Ethical Pharmaceuticals of Merit since 1878

### THEORATE

Robins

COR VITALIS  
FACILITATIS  
COMES ET  
ARTE PRIN.





# life-long constipation

due to bulk deficiency

## corrected in days

*YEARS OF OBSTINATE CONSTIPATION* can be corrected in days with Cellothyl. Barger<sup>1</sup> reports "striking" results in patients with "no ordinary form of constipation" but with life-long dysfunction.

IN "*ORDINARY*" CASES, Cellothyl is equally effective. In patients taken at random from routine office practice, results were "excellent" or "good" in 92% of cases.<sup>2</sup>

*PHYSIOLOGIC CORRECTION—IN THE COLON:* Cellothyl provides bulk where it is needed—in the colon. It passes through the stomach and upper intestines as a fluid and thickens to a smooth gel in the colon to provide bulk for soft, moist, easily passed stools.

The usual starting dose is 3 tablets t.i.d., each dose accompanied by at least one glass of water. Daily fluid intake should be high. As normal function returns, the Cellothyl dosage may be gradually reduced.

1. *Gastroenterology* 13: 275 (Oct.) 1949. 2. *N. Y. State J. Med.* 48: 1822 (Aug.) 1948.

  
 CELLOTHYL TABLETS

+

  
 WATER

+

  
 TIME

=

  
 PHYSIOLOGIC COLONIC BULK

# Cellothyl<sup>®</sup>


brand of methylcellulose  
especially prepared



**CHILCOTT**  
*Laboratories*

DIVISION OF The Maltine Company

MORRIS PLAINS, NEW JERSEY



*Because*

*patients can't*

**"SLEEP OFF" hypertension...**

*prolonged vasodilation should accompany sleep  
as well as the day's activities. (One more reason why*

*NITRANITOL is the most universally prescribed  
drug in the management of hypertension.)*

# NITRANITOL®

FOR GRADUAL, PROLONGED, SAFE VASODILATION



**When vasodilation alone is indicated.** *Nitranitol.* ( $\frac{1}{2}$  gr. mannitol hexantrate.)

**When sedation is desired.** *Nitranitol with Phenobarbital.* ( $\frac{1}{2}$  gr. Phenobarbital combined with  $\frac{1}{2}$  gr. mannitol hexantrate.)

**For extra protection against hazards of capillary fragility.** *Nitranitol with Phenobarbital and Rutin.* (Combines Rutin 20 mg. with above formula.)

**When the threat of cardiac failure exists.** *Nitranitol with Phenobarbital and Theophylline.* ( $\frac{1}{2}$  gr. mannitol hexantrate combined with  $\frac{1}{2}$  gr. Phenobarbital and 1½ grs. Theophylline.)

# MODERN MEDICINE

THE JOURNAL OF DIAGNOSIS AND TREATMENT

## EDITORIAL BOARD

James T. Case, M.D., *Chicago*  
Franklin D. Dickson, M.D., *Kansas City*  
Julius Hess, M.D., *Chicago*  
Walter B. Hoover, M.D., *Boston*  
Foster Kennedy, M.D., *New York City*  
John C. Krantz, Jr., PH.D., *Baltimore*  
A. J. Lanza, M.D., *New York City*  
Milton S. Lewis, M.D., *Nashville*  
George R. Livermore, M.D., *Memphis*  
Francis W. Lynch, M.D., *St. Paul*  
Cyril M. MacBryde, M.D., *St. Louis*  
Karl A. Meyer, M.D., *Chicago*  
Alton Ochsner, M.D., *New Orleans*  
Robert F. Patterson, M.D., *Knoxville*  
Edwin B. Plimpton, M.D., *Los Angeles*  
Fred W. Rankin, M.D., *Lexington, Ky.*  
John Alton Reed, M.D., *Washington*  
Rufus S. Reeves, M.D., *Philadelphia*  
Dalton K. Rose, M.D., *St. Louis*  
Howard A. Rusk, M.D., *New York City*  
Roger S. Siddall, M.D., *Detroit*  
James S. Simmons, M.D., *Boston*  
W. Calhoun Stirling, M.D., *Washington*  
Frank P. Strickler, M.D., *Louisville*  
Richard Torpin, M.D., *Augusta, Ga.*  
Paul M. Wood, M.D., *New York City*

---

SCIENCE WRITERS: Thomas B. Gibbons, M.D.,  
Norman Shrifter, M.D., T. C. Paper-  
master, M.D., H. D. Pocock, M.D., Ber-  
nardine Lufkin, Elizabeth Kane, Shanna  
McGee


TRANSLATOR: John W. Brennan, M.D.

## EDITORIAL CONSULTANTS

Carolyn G. Adams, M.D.  
E. R. Anderson, M.D.  
Joe W. Baird, M.D.  
Edward P. Burch, M.D.  
James B. Carey, M.D.  
C. D. Creevy, M.D.  
C. J. Ehrenberg, M.D.  
W. K. Haven, M.D.  
Charles Hymes, M.D.  
Miland E. Knapp, M.D.  
Ralph T. Knight, M.D.  
John F. Pohl, M.D.  
Leo Rigler, M.D.  
Wallace P. Ritchie, M.D.  
M. B. Sinykin, M.D.  
A. V. Stoesser, M.D.  
Arthur L. H. Street, LL.B.  
Marvin Sukov, M.D.  
Harry A. Wilmer, M.D.

## EDITORIAL STAFF

A. E. Hedback, M.D.  
*Editor*  
Thomas Ziskin, M.D.  
*Associate Editor*  
Maurice B. Visscher, M.D.  
*Consultant in Sciences*  
Mark S. Parker  
*Executive Editor*  
Sarah A. Davidson  
*Managing Editor*  
Daisy Stilwell  
*Art Editor*  
James Niess  
*Editorial Board  
Secretary*  
Jeanne E. Feldmeir  
Diane N. Murray  
*Editorial Assistants*




MULTI-PURPOSE TABLE MODEL A, TYPE 1

*with a Ritter*  
**MULTI-PURPOSE TABLE**

**W**HATEVER the problem, you can always examine patients at the height most convenient for you with a Ritter Multi-Purpose Table. This motor-driven table gives rapid, smooth adjustment to any required height by slight toe pressure on the foot pedal . . . your assurance of always working at *your comfort level*.

MULTI-PURPOSE TABLE  
MODEL A, TYPE 2



*If you're vacationing in the East, be sure to visit our plant.*

The Model A, Type 1, shown above, adjusts to any position from full horizontal to chair. Rotates 180° with range of elevations, 27" to 45" or 23" to 41", from table top to floor.

The Model A, Type 2 Multi-Purpose Table, shown at left, has the same features as the Type 1, plus a special adjustable knee rest for proctological work. Extreme tilt 55°. Elevation range 31" to 49", from table top to floor.

See your surgical dealer today for a demonstration of these Ritter Multi-Purpose Tables.





**drowsiness** *minimized . . .*  
*allergic patients remain alert . . .*

Clinical reports describing the use of Thephorin in 2564 patients with hay fever and other allergies indicate an incidence of drowsiness of only 2.92%. In contrast with other antihistamines, Thephorin can therefore be given to motorists and other patients who have to remain alert. Highly effective and well tolerated in most cases,

Thephorin is available in 25-mg tablets and as a palatable syrup which permits convenient adjustment of dosage.

HOFFMANN-LA ROCHE INC • NUTLEY 10 • N. J.

**Thephorin®**  
*brand of phenindamine*

**'Roche'**

LETTER FROM THE EDITOR.....	14
CORRESPONDENCE.....	18
QUESTIONS & ANSWERS.....	24
FORENSIC MEDICINE.....	30
SPECIAL ARTICLE	

Physical Medicine and Rehabilitation in Poliomyelitis <i>Robert L. Bennett</i> .....	43
--	----

#### MEDICINE

Adductor Longus Syndrome <i>Janet Travell</i> .....	48
Treatment of Venous Thrombosis <i>F. Richard Pierce and Trento J. Domenici</i> .....	49
Current Histamine Therapy <i>Lester S. Blumenthal</i> .....	51
Labial Diphtheria.....	53
Aerobic Fecal Bacteria.....	53
Diagnosis of Addison's Disease.....	54
Liver Extract Injections.....	54
Absolute Lymphocytopenia.....	54

#### SURGERY

Commissurotomy for Mitral Stenosis <i>Robert P. Glover, Thomas J. E. O'Neill, and Charles P. Bailey</i> .....	55
Respiratory Borborygmus.....	56
Thymectomy for Myasthenia Gravis <i>Henry R. Viets</i> .....	57
Improved Biliary T-Tube <i>R. Russell Best</i> .....	58
Ulcerative Colitis <i>J. M. Rice-Oxley and Sidney Truelove</i> .....	59



## Contents for August 1 1950

MODERN MEDICINE  
VOL. 18, NO. 15

THE MAN ON THE COVER is Robert L. Bennett, M.D., a specialist in physical medicine whose particular interest is poliomyelitis. Dr. Bennett is Director of Physical Medicine at the Georgia Warm Springs Foundation, which was founded by the late President Franklin D. Roosevelt. Dr. Bennett is a frequent contributor of articles on physical medicine to scientific journals and is author of the Special Article, "Physical Medicine and Rehabilitation in Poliomyelitis," on page 43 of this issue.





*Contents*  
*for*  
*August 1*  
*1950*

CONTINUED

ORTHOPEDICS

- Elastic Hand Splints  
*Sterling Bunnell and*  
*Lot D. Howard, Jr.*..... 62
- Pulp Space Infection of the Finger  
*Harold Bolton, P. J. Fowler,*  
*and R. P. Jepsen*..... 64

OBSTETRICS

- Frequency of Plural Births..... 65

PEDIATRICS

- Parenteral Fluid Therapy for Children  
*Albert M. Hand and*  
*C. R. Leininger*..... 66

LARYNGOLOGY

- Perforation of the Esophagus  
*John D. Kernan*..... 68

OPHTHALMOLOGY

- Diabetic Retinopathy  
*Bruce L. Kantar*..... 69

SURGICAL TECHNIGRAM

- Femoral Hernia  
*F. M. Al Akl*..... 70

MEDICAL FORUM

- Non-Union of the Medial Malleolus..... 76

DIAGNOSTIX..... 78

SHORT REPORTS..... 88

WASHINGTON LETTER..... 94

CURRENT BOOKS & PAMPHLETS..... 106

PATIENTS I HAVE MET..... 110

MODERN MEDICINE, The Journal of Medical Progress, is published twice monthly on the first and fifteenth of each month at Minneapolis, Minn. Subscription rate: \$5.00 a year, 25c a copy. Business Manager: M. E. Hertz. Address editorial correspondence to 84 South 10th Street, Minneapolis 3, Minn. Telephone: Bridgeport 1291. ADVERTISING REPRESENTATIVES: New York 17: Lee Klemmer, George Doyle, Bernard A. Smiler, 50 East 42nd Street, Suite 401. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Hertz, 20 North Wacker Drive, Suite 1921. Telephone: Central 6-4619. SAN FRANCISCO 4: Duncan A. Scott & Co., Mills Bldg. Telephone: Garfield 1-7950. LOS ANGELES 5: Duncan A. Scott & Co., 2979 Wilshire Blvd. Telephone: Dunkirk 8-4151.





## **Stubborn TRICHOMONIASIS**

# **yields to ARGYPULVIS in 98% OF CASES\***

\*Report of Reich, Button and Nechtow in *Surgery, Gynecology and Obstetrics*... (See coupon offer.)

**T**his newest adaptation of one of medicine's standbys—ARGYROL—gives the physician a most effective and simplified therapy for better control of trichomoniasis. The same effective

powder used in office treatments is available for supplementary home use to speed results. For better acquaintance with the ARGYPULVIS technique, use the coupon below.

### **....in two convenient forms .....**



**For Use by the Physician.** 7-gram bottles fitting Holms-spray or equivalent powder-blower (in cartons of 3).

**For Home Use by the Patient.** 2-gram capsule for insertion by the patient (in bottles of 12)



**INTRODUCTORY TO PHYSICIANS:** \*On request we will send professional samples of ARGYPULVIS (both forms), together with a reprint of the Reich, Button and Nechtow report. (Use coupon.)

A. C. Barnes Company  
Dept. MM-80, New Brunswick, N. J.

Name.....

Address.....

City.....State.....

## **ARGYPULVIS**

ARGYROL and ARGYPULVIS are registered trademarks, the property of  
**A. C. BARNES CO., NEW BRUNSWICK, N. J.**

## LETTER FROM THE EDITOR

*Dear Reader:*

Two plus two does not always equal four.

Perhaps you remember the story of the fullback who was better at plunging through the line than pulling down a D. He had failed in every test the math instructor had given. The coach pleaded for one more chance. The instructor wearily agreed that if the gridder could answer one question correctly the minimum eligibility requirements would be met. The question: What does two plus two equal?


The fullback chewed his pencil long and thoughtfully. Then his eyes brightened and he triumphantly wrote "three." The instructor left the room without a word. The coach, on the verge of apoplexy, turned to the player.

"Yah big lunk," he exploded, "everybody knows the answer is five."

The equations are not that simple in medicine, but opinion as to the correct answer is often just as diverse. This diversity of opinion makes the Medical Forum one of the most interesting and stimulating departments in *Modern Medicine*.

As each issue is being prepared, our Editorial Committee selects for discussion two or three reports that touch on controversial points. Readers are encouraged to express their opinions. Often proofs are sent for comment to practitioners known to be particularly interested in the diagnostic or therapeutic problem involved. Usually the majority agree that the answer is four, but on occasion wide disagreement is revealed.

The Medical Forum is a sounding board for the serious and considered opinion of every doctor. He may be famed throughout the nation or he may be unknown outside his bailiwick. If he has a significant comment to make, the Medical Forum is available to him.



EDITOR

three

wise



## considerations

In biliary stasis, Caroid and Bile Salts Tablets fulfill the need for a desirable threefold action as a—

- 1...**choleretic** to produce increased bile flow
- 2...**digestant** to assist digestion
- 3...**laxative** to induce peristaltic action and help reestablish normal function.

Caroid and Bile Salts Tablets have gained an enviable reputation—over a period of years—as an effective aid in the treatment of biliary stasis, cholecystitis, constipation of biliary origin.

Available in bottles of 20, 50, 100, 500 and 1000.  
tablets of...

## Caroid and Bile Salts

with Phenolphthalein

American  
Ferment

trial supply on request

Company, inc. 1450 Broadway, New York 18, N. Y.

*Announcing...*

# 'PERAZIL' BRAND

CHLORCYCLIZINE HYDROCHLORIDE

*a (new type) antihistaminic*

*Relief from allergic symptoms for  
12 to 24 hours with a single dose*



'Perazil' brand Chlorcyclizine Hydrochloride is a completely new type of antihistaminic, its distinctive component being a piperazine ring instead of the usual ethylenediamine grouping. This uniquely different chemical structure results in a prolongation of action—up to 24 hours following a single 50 mg. dose.<sup>1</sup> In contrast to many other antihistaminic compounds, 'Perazil' exhibits a low incidence of side-effects despite its high potency and prolonged effectiveness.

**INDICATIONS:** Hay fever, vasomotor rhinitis, urticaria, allergic dermatitis and pollen asthma.


**DOSAGE:** 50 mg. (one product) once daily with water; may be increased to two or three times daily if required in very severe cases.

**PREPARATION:** 'Perazil' brand Chlorcyclizine Hydrochloride 50 mg. Compressed (scored). Bottles of 100.

1. Jares, S. H.: *Annals of Allergy*, Vol. 7, No. 4 (July-Aug.) 1949



**BURROUGHS WELLCOME & CO. (U. S. A.) INC., TUCKAHOE 7, N. Y.**



when your patient sees it daily . . . his "tonic" becomes an ever-present symbol of the reassuring and comforting fact that he is "in the care of his physician". Physicians know that, in addition to its tone-restoring and appetite-stimulating effects, this psychological aspect of a good tonic can often produce striking results. **NEURO PHOSPHATES** and **THERANATES** are both available in 12 fl. oz. bottles.

*Smith, Kline & French Laboratories, Philadelphia*

## **Eskay's Neuro Phosphates\***

*a palatable and effective tonic*

## **Eskay's Theranates\***

*the formula of famous Neuro Phosphates, plus Vitamin B<sub>1</sub>*

\*T. M. Reg. U. S. Pat. Off.

# Correspondence

Communications from the readers of MODERN MEDICINE are always welcome. Address communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.

## How to Advise Psychiatry?

TO THE EDITORS: Occasions arise when a general practitioner has reason to suggest a psychiatric consultation to a patient but is reluctant to do so. There is the fear that the patient will be incensed because he thinks that the physician is implying doubt as to his sanity, no matter the reason for the consultation. The situation is a delicate one, to be handled with tact and discrimination.

I would be very interested in the experiences of others with this problem and in receiving suggestions as to how this matter may best be handled.

EDWARD DENGROVE, M.D.  
Asbury Park, N. J.

## Cause of Sexual Impotence

TO THE EDITORS: In the Questions & Answers section of a recent issue of your excellent journal (*Modern Medicine*, June 1, 1950, p. 28) is a brief discussion of the etiology and therapy of loss of sustained erection. The same topic is dealt with in Queries and Minor Notes of the June 3 issue of the *Journal of the American Medical Association*.

In each instance the consultant pointed out that usually such cases are on a psychogenic, hormonal, or neurogenic basis. I should like to

emphasize that insidious thrombosis of the terminal aorta is occasionally responsible for sexual impotency.

Slowly progressive thrombosis of the terminal aorta should no longer be considered a disease of great rarity. The syndrome is entirely unrelated to and is to be sharply differentiated from the acute embolic obstruction and rapidly progressive thrombosis so commonly seen in cardiac patients.

The recent marked increase in the number of reported cases suggests that the disease, so-called Leriche's syndrome, occurs with much greater frequency than is generally appreciated. Elkin and Cooper (*Ann. Surg.* 130:417, 1949) have recently reviewed the clinical features and treatment of insidious thrombosis of the terminal aorta. These authors discovered 10 cases in twenty months. I have had 2 personal cases during the past four months.

A provisional diagnosis of thrombotic occlusion of the distal aorta may be made whenever a patient complains of exertional pain in the hips and thighs, easy fatigability of the legs, or sexual impotency and examination discloses no pulsations in vessels distal to the umbilicus. Aortography will confirm the diagnosis and reveal the level of obstruction.

MANY DOCTORS THROUGHOUT THE NATION  
REPORT BETTER RESULTS WITH

*THE*  
*Stuart Formula Liquid*

**COMPARE**  
**COMPLETENESS**

An aqueous preparation containing vitamins A & D,  
*plus* a complete B Complex including natural B Complex factors  
from malt and rice polish concentrate *plus* minerals.

**TASTE**

There is no better tasting liquid multivitamin. It really tastes *good*.

**COST**

1 pint costs your patients **\*2.50**

the **Stuart**  
Company

the **Stuart**  
**formula**  
**liquid**

Vitamins  
A D B<sub>1</sub> B<sub>2</sub> P-P B<sub>6</sub> E  
*Pantherol*  
including entire  
B Complex  
Minerals  
Malt

the **Stuart**  
Company

NET CONTENTS, ONE PINT

# EXTRA *Protection*

## AT NO EXTRA COST TO PATIENTS

THE STUART LYOPHILATED\* MULTIVITAMIN DROPS is a product which stays fresh...yet requires no expiration date, no refrigeration, and no addition of alcohol...in other words, a multivitamin drop upon which you can really depend. The Stuart method of lyophilizing the B Complex factors and C in one bottle—and separating the aqueous dispersion of Vitamins A and D in another bottle—assures full label potencies.

Low in cost to patients, 30 cc. **"2.35"**





Treatment consists in bilateral lumbar sympathectomy, which destroys the vasoconstrictor nerve supply to the collateral vessels distal to the obstruction. If treatment is instituted before gangrene has appeared, relief of symptoms is remarkable in most cases.

It is to be emphasized that the thrombosis develops very gradually; the average duration of symptoms in these cases of Elkin and Cooper was seven years. The disease is not confined to the elderly arteriosclerotic group. Leriche's youngest patient was a woman of twenty-nine.

Thrombotic obliteration of the terminal aorta should be considered in the differential diagnosis in cases of sexual impotency.

WALTER F. BECKER, M.D.

Little Rock, Ark.

#### Annual Arrival

TO THE EDITORS: The long-awaited *Annual* has been received, and your organization has done it again—meeting all the claims you had made and meriting a thunderous outburst of huzzahs!

HAROLD T. LOKSA, M.D.

Bayonne, N. J.



# PRURITUS

...due to Insect Bites  
by Poisoning •  
Localized Vesicular Areas



## CALAMATUM

(NASON'S)


affords immediate relief for the itching and discomfort of skin affections prevalent during the summer months. It is a cream embodying Calamine with Zinc Oxide and Campho-Phenol in a non-greasy base. CALAMATUM dries at once, adhering to the lesion and thus localizing the infection by preventing spread of any exudate. By alleviating itching with consequent desire for relief by scratching, it reduces the dangers of secondary infection.

### WON'T RUB OFF

Easy application without messy liquids and embarrassing bandages, and the handy tube instead of a fragile bottle of lotion encourage applications at any time. In 2-oz. tubes at druggist or direct.

TAILBY-NASON COMPANY

Kendall Sq. Station • BOSTON 42, MASS.





combined anesthetic-lubricant  
action facilitates cystoscopy, proctoscopy  
and other endoscopic procedures.



relieves anal and rectal pain  
especially valuable postoperatively.  
Each package contains an anorectal applicator.

protective anesthetic dressing for  
painful first and second degree burns.

# Questions & Answers

All questions received will be answered by letter directed to the petitioner; questions chosen for publication will appear with the physician's name deleted. Address all inquiries to the Editorial Department, MODERN MEDICINE, 84 South Tenth Street, Minneapolis 3, Minnesota.

**QUESTION:** Please outline a recommended course of therapy for a patient whose prostate gland is infected with *Trichomonas vaginalis*.

M.D., West Virginia

**ANSWER:** By Consultant in Urology. Eradication of *Trichomonas vaginalis* from so complicated a structure as the prostate gland is very difficult without an effective agent that will kill the organism upon systemic administration. Since something like 100 different drugs and chemicals have been recommended for treatment of this disease in the female, we may conclude that no one method is likely to be satisfactory.

The first step is to make sure that the patient does not have urethral stricture and that no complicating bacteria are associated with the infection. This is best done by culturing the combined urine and prostatic secretion. If bacteria are found, appropriate antibiotic therapy should be utilized. If stricture is present, urethral dilatation should be employed. If pus is found in the prostatic secretion, massage two or three times weekly may be desirable.

The various arsenicals used in the treatment of syphilis may be tried, either orally or intravenously ac-

cording to the preparation, but without much prospect of benefit. The instillation into the prostatic and anterior urethra of a 3% solution of neosarsphenamine may also be tried.

**QUESTION:** Why does neurologic examination of a sixty-five-year-old man with cerebral hemorrhage resulting in complete hemiplegia of the right side yield a positive Babinski on the unaffected left foot and a normal plantar response on the paralyzed right side?

M.D., Massachusetts

**ANSWER:** By Consultant in Neurology. The patient described undoubtedly has a very extensive lesion with a great deal of irritation of the brain bilaterally from the edema. It is not uncommon in such cases to obtain a positive Babinski sign bilaterally or on the side opposite the lesion.

**QUESTION:** Please tell me the simplest practical way to determine by roentgenography if the bony pelvis is large enough to permit normal delivery at term.

M.D., Tennessee

**ANSWER:** By Consultant in Roentgenology. A number of methods of pelvic mensuration by roentgen study

# *For Greater Variety in the* **DIABETIC DIET**


Experience shows that the use of Knox Unflavored Gelatine in the diabetic dietary is a valuable help to the patient, in four ways:

1. It allows greater freedom in the choice of foods,
2. Permits dishes containing a minimum of available carbohydrates,
3. Makes restricted portions as appetizing as those of the normal diet,
4. In diabetic reducing diets, it makes possible a wide variety of dishes that are high in residue while low in calories.

Knox Gelatine, unlike factory-flavored gelatin dessert powders, with their high sugar and acid content, is all protein, no sugar.

## **FREE** DIABETIC DIET GUIDE

This 48-page book contains "diabetic arithmetic," food composition charts, the various standards of diabetic nutrition, and 78 tested diabetic menus and recipes, each with a chart of carbohydrate, protein, fat and caloric content. Of high value to diabetic patients, a supply of these books is free upon request. Write Knox Gelatine, Box R-35, Johnstown, N. Y.



Available at  
grocery stores  
in condensed  
4-envelope and  
38-envelope  
packages.



## QUESTIONS & ANSWERS

are described by H. C. Moloy and P. C. Swenson in Chapter 10 of the looseleaf textbook, *Diagnostic Roentgenology*, edited by Ross Golden and published by Thomas Nelson & Sons. Additional references are: *Clinical Roentgenology of Pregnancy* by William Snow, Charles C Thomas, 1942; "The Accuracy of Roentgen Estimates of Pelvic and Fetal Diameters," A. L. Dippel and E. Delfs, *Surg., Gynec. & Obst.* 72:915-922, 1941; "Roentgen Pelvimetry and Fetometry," P. C. Hodges, *Am. J. Roentgenol.* 37:644-662, 1937; "Elastic Ruler for Roentgen Pelvimetry," Cesare Gianturco, *Radiology* 49:95-96, 1947.

Probably the simplest practical method for determining mensuration


of the bony pelvis is that of Snow. A slide rule called Snow Calculator is available for easy computation. Directions are given with the rule. By making two films, an antero-posterior and a lateral, distortion can be corrected and a fairly accurate measurement of the pelvic diameters determined. The methods of Dippel or Hodges, which are quite similar, are probably more precise but require more complicated calculations than the Snow method.

**QUESTION:** Can large doses of streptomycin or aureomycin cause swelling of the liver?

M.D., New York

**ANSWER:** By Consultant in Internal Medicine. No.

**ARE YOUR  
COLLECTIONS  
SLIPPING**



**Collectvelope**

**A NEW COLLECTION IDEA THAT  
OBTAINS REMARKABLE RESULTS**

"Collectvelope" is a COMBINATION request for payment and reply envelope. It is breaking all doctors' collection records because it makes it easy for patients to pay.

**SEE THIS AMAZING NEW ITEM  
ACTUAL SAMPLES FREE ON REQUEST**

• Reg. U.S. Pat. Off.

**DOLLARS COLLECTED ARE DOLLARS EARNED**

PROFESSIONAL PRINTING CO., INC.  
202 Tillary Street, Brooklyn 1, N. Y. 2-8-0  
Gentlemen: Send me actual samples and  
all details on the NEW "Collectvelope."  
Dr. \_\_\_\_\_

**AS NEW AS TOMORROW**



**PROFESSIONAL  
PRINTING COMPANY, INC.**

202-204 TILLY ST. BROOKLYN 1, N. Y.



***depression*** causes ...



***loss of appetite,***  
which leads to ...



***inadequate dietary intake***  
and, consequently, to ...



***B vitamin deficiency,***  
resulting finally in ...



***apathy, fatigue and***  
***physical debility***

**to arrest the "downward spiral"**

'Dexedrine'\* plus essential B vitamins

**Theptine**

**an antidepressant and nutrient elixir**

Available in 12 fl. oz. bottles.

Smith, Kline & French Laboratories • Philadelphia

\*'Theptine' and 'Dexedrine' are S.K.F. Trademarks

# RIASOL in



**BEFORE USING RIASOL**

## **THOUSANDS PRESCRIBE RIASOL**

RIASOL CONTAINS 0.45% MERCURY CHEMICALLY COMBINED WITH SOAPS, 0.5% PHENOL AND 0.75% CRESOL IN A WASHABLE, NON-STAINING, ODORLESS VEHICLE

**APPLIED LOCALLY—SIMPLE TO USE**



# PSORIASIS

---



**AFTER USING RIASOL**

**SEND FOR A CLINICAL PACKAGE**

**PROVE RIASOL YOURSELF**

**SHIELD LABORATORIES**

**12860 MANFIELD AVENUE, DETROIT 27, MICHIGAN**

# Forensic Medicine

COMPILED BY ARTHUR L. H. STREET, LL.B.

**PROBLEM:** In the trial of a personal injury suit could the jury determine, as a matter of inference, that plaintiff's injury would be permanent and involve future suffering, when roentgenograms and medical testimony disclosed a progressive physical condition, which in the light of common experience warranted that inference?

**COURT'S ANSWER:** Yes.

The Oklahoma Supreme Court decided that the evidence disclosed both objective and subjective conditions, involving progressive traumatic arthritis and spinal injury (212 Pac. 2d 469).

**PROBLEM:** An indictment charged accused with [1] practicing medicine without a license, [2] unlawfully prefixing "Dr." to his name, and [3] using the abbreviation in an occupation relating to public health and diagnosis and treatment of disease. Could a conviction on the second count alone stand?

**COURT'S ANSWER:** No.

The Florida Supreme Court decided that the conviction implied an acquittal on the charges of practicing without a license and using "Dr." in connection with medical practice. (The court noted that many "doctors" are not medical doctors and that there was no proof that, if defendant did use the title, "Dr.," he intended to pose as a medical practitioner (45 So. 2d 118).

**PROBLEM:** The jury in a bastardy proceeding found that respondent was the father of twins, although careful blood-grouping tests ordered by the court indicated nonpaternity. Was he entitled to a new trial?

**COURT'S ANSWER:** Yes.

Eleven blood tests were made by a widely recognized expert. The blood was secured by 3 physicians, who testified in detail as to the steps taken. The results were identical in each test, showing that the blood of man, woman, and twins all belonged to Group A. However, the man's blood was Type N, whereas that of the mother and one twin was Type M and that of the other twin was Type MN.

The expert based his opinion that the man could not be the father on two grounds: [1] the biologic rule that a parent with Type N blood cannot have a child with Type M, and [2] the theory that the father of twins must be the same man.

Upholding a verdict of nonpaternity, the Maine Supreme Judicial Court found it unnecessary to consider the question as to whether there must be an identical father of twins, because the verdict was indivisible; if the evidence excluded paternity as to one child there must be a new trial.

The court reaffirmed what it had said in a previous case, that the

(Continued on page 34)

# quick relief and healing

in  
peptic  
ulcer...

**Resinat**, pepsin inactivator and antacid, brings quick relief of pain and speeds healing of peptic ulcer.

Weiss, S., et al.,<sup>1</sup> used Resinat in the treatment of 120 ulcer patients. These investigators report "symptomatic relief occurred within 48-72 hours and x-ray follow-up showed regression of ulcer crater in two to four weeks."

**Resinat** is insoluble, chemically and physiologically inert. It does not remove chlorides, phosphates, vitamins or minerals from the body. It does not alkalize the system or cause acid rebound.

Resinat inactivates pepsin and neutralizes excess gastric acidity.


Available in Capsules, 0.25 Gm. — Tablets, 0.5 Gm. — Powder, 1 Gm. Packets.


1. Weiss, S., et al.: *Rev. Gastroenterology* 16:501-509 (June) 1949. Literature and samples available.

*More than  
Half a Century  
of Service  
to the Medical Profession*



**The National Drug Company**  
*Philadelphia 44, Pa.*





## Picture the patient ... *Routine procedure*

Here are instruments ideally suited to the widening demands of medical photography.

Exactly framed, needle-sharp pictures are routine with the new Kodak Flurolite Camera Combination and the Kodak Flurolite Enlarger assembled for photographing gross specimens . . . thanks to the revolving swing back, the easy adjustments, the long bellows draw.




The Camera Combination uses  $2\frac{1}{4}$  x  $3\frac{1}{4}$ -inch sheet film . . . can easily

be converted to accept 35mm roll film. Interchangeable color-corrected Kodak Ektar Lenses in Flash Supermatic Shutters assure brilliant results—black-and-white or full-color . . . flood or flash.

*Additional uses of the Kodak Flurolite Enlarger (with accessories):* Copying, slide-making, cine-titling, photomicrography, and microfilming.

See the new Kodak Flurolite Camera Combination and Kodak Flurolite Enlarger at your Kodak dealer's . . . or write for further information. Eastman Kodak Company, Medical Division, Rochester 4, N. Y.

*Serving medical progress*




Reproductions A and B show gross aspects of diseased lung after pneumonectomy. Reproduction C shows photomicrograph made from histological section of affected lung. Reproduction D shows setup of Kodak Fluorolite Enlarger fitted with Camera components for the photography of gross specimens.

## *with more and more physicians*

*Kodak products for the medical profession include:*

X-ray films; x-ray intensifying screens; x-ray processing chemicals; electrocardiographic papers and film; cameras—still- and motion-picture; projectors—still- and motion-picture; enlargers and printers; photographic film—color and black-and-white (including infrared); photographic papers; photographic processing chemicals; synthetic organic chemicals; microfilming equipment and microfilm.



*Kodak Fluorolite Camera Combination  
with Kodak 35mm Film Adapter A.*

*rough Photography and Radiography*

**Kodak**  
TRADE-MARK

## FORENSIC MEDICINE

statute authorizing use of blood tests in bastardy proceedings accepts the scientific view "that even though such tests cannot prove paternity, they may in certain instances disprove it. . . . Such a law goes beyond the opinion of an expert. The jury has the duty to determine if the conditions existed which made the biological law operative. That is, . . . were the tests properly made? If so made, the exclusion of the respondent as father of one child follows irresistibly."

The court added that its decision in the previous case did not indicate "that a jury may give such weight as it may desire to biological law." If the jury found in this case that the results of the tests were inac-

curate, it must have been on mere conjecture or "understandable sympathy for the mother."

But the court rejected argument on behalf of respondent that the court had no jurisdiction to entertain a bastardy proceeding involving twins—that there should have been a separate proceeding as to each child (69 Atl. 2d 670).

**PROBLEM:** Was a physician guilty of murder in the second degree if he intentionally used means to destroy an unborn child or to produce miscarriage, without intending, in good faith, to save the mother or the child?

**COURT'S ANSWER:** Yes.

The West Virginia Supreme Court of Appeals upheld conviction in

For controlled  
Rate and Rhythm..



\*Not an advertisement  
mixture of glycosides



Through more precise control of contractile force and rhythm, Digitaline Nativele maintains the maximum efficiency obtainable. Maintenance is positive because absorption is complete and the rate of dissipation is uniform; providing full digitalis effect between doses.

# DIGITALINE NATIVELE

Chief active principle\* of digitalis purpurea

(digitoxin)

**MAINTENANCE:** 0.1 or 0.2 mg. daily depending on patients' response.

**CHANGE-OVER:** 0.1 or 0.2 mg. Digitaline Nativele replaces 0.1 or 0.2 gm. whole leaf.

**RAPID DIGITALIZATION:** 0.6 mg. initially, followed by 0.2 or 0.4 mg. every 3 hours until digitalized.

Send for brochure "Modern Digitalis Therapy" Varick Pharmacal Co. Inc. (Div. E. Fougere & Co. Inc.), 75 Varick St., New York

• *complete-vitamin formula . . .*

• *water miscible . . .*

• *non-alcoholic . . .*

• *inexpensive . . .*



*White's*

## Multi-vi drops

Each 0.6 cc. contains:

Vitamin A 5000 U.S.P. units • Vitamin D<sub>3</sub> 1000 U.S.P. units  
Thiamine Hydrochloride 1.0 milligram • Riboflavin 0.4 milligram  
Pyridoxine Hydrochloride 1.0 milligram • Panthenol\* 2.0 milligrams  
Nicotinamide 10.0 milligrams • Ascorbic Acid 50.0 milligrams

\*a specially-prepared, more stable analogue of pantothenic acid.

White Laboratories, Inc., Pharmaceutical Manufacturers, Newark 7, N. J.



## FORENSIC MEDICINE

this case. The defendant was sentenced to imprisonment for not less than five years and not more than eighteen.

Under a statute of that state, the court decided that an indictment charging murder was proper without specifying that the fatal act consisted in performing a criminal abortion.

The court also said that the trial judge did not err in permitting the prosecution to show that accused had performed other criminal abortions within five years—not as tending to prove that the accused had committed the specific offense charged, but as indicating a systematic course of criminal conduct (57 S.E. 2d 513).

**PROBLEM:** Was a family physician, who remained in charge of the case, jointly liable with an orthopedist for negligent diagnosis and treatment of a fractured humerus?

**COURT'S ANSWER:** Yes.

The Wisconsin Supreme Court stressed the facts that the family physician had remained in active charge and was negligent in not securing roentgenograms to determine the position of the bones that had been set and the progress of healing. These facts, the court said, differentiated the case from an earlier one (183 Wis. 382, 197 N.W. 333) in which a family physician was held not jointly responsible with an operating surgeon, whom he assisted (41 N.W. 2d 620).

**For Finger Freedom You'll Enjoy**

Specify  
Specify  
Specify

**ROLLPRUF**

**PIONEER**  
SURGICAL GLOVES



You do two things when you specify Pioneer Rollprufs. You get longer surgical glove wear and economy — and you give yourself a glove you enjoy wearing.

### **Here's why:**

**FLAT BANDED CUFFS** — exclusive with Rollprufs. Wrists won't roll down during surgery — reduce tearing.

**COMFORT-FIT** — all Rollprufs, latex and neoprene, are more comfortable, less tiring in long wear.

**PIONEER ROLLPRUFS** — Natural latex and DuPont neoprene. Neoprene Rollprufs in the new hospital green color for easy sorting — are free of the dermatitis inducing allergen sometimes found in natural rubber.

Rollprufs are more for your money. Processed to stand extra sterilizings, tough yet sheer, they afford added sensitivity to surgeon's fingers. SPECIFY Rollprufs from your supplier or write us.

The Pioneer Rubber Company • 751 Tiffin Road • Willard, Ohio



**A No. 1  
DRUG  
for the  
No. 1  
ALLERGY**

# **Pyribenzamine®**

*for Hay Fever*

More PYRIBENZAMINE than any other antihistaminic has been prescribed for hay fever and other allergies. Wide clinical use has established that PYRIBENZAMINE (tripelennamine) hydrochloride provides maximum allergic relief with minimal side effects.

**Ciba** PHARMACEUTICAL PRODUCTS, INC., SUMMIT, N. J.


2/1618M

There is an important difference in Elastic Bandages!

# TENSOR\*

ELASTIC BANDAGE

is woven with *Live Rubber Thread!*



And *what* a difference that makes! Since TENSOR is woven with *live rubber thread*, it offers you these unequalled advantages over rubberless bandages.

## TENSOR:

- exerts uniform, controlled pressure without binding.
- has constant elasticity—you don't find it in rubberless bandages!
- stays put—needs no frequent readjustments—is comfortable to wear.
- is inconspicuous—women patients will wear it.
- can be laundered repeatedly without loss of elasticity.

**Ideal For Every Elastic Bandage Use!**

For your patients' support and comfort,  
prescribe Bauer & Black Elastic Supports

**BAUER & BLACK ELASTIC STOCKINGS**... 2-way stretch, easy to fit. Women appreciate their neutral and inconspicuous color.



**BAUER & BLACK SUSPENSORIES**... as a palliative and for treatment of diseases of the scrotum.



\*Reg. U. S. Pat. Off.

Products of

**BAUER & BLACK**


Division of The Kendall Company, 2500 S. Dearborn St., Chicago, 16

**FIRST IN ELASTIC SUPPORTS**

# medical management that actually promotes faster, better HEALING in PEPTIC ULCER

*Not just another  
antacid or  
prokinetic...*

but a new and positive  
approach to peptic-ulcer  
therapy. Each capsule  
provides, for the first  
time, the complementary  
action of:



**ALLANTOIN**  
60 mg.

Minimizes gastric  
inflammation and  
stimulates healing

**ASCORBIC ACID**  
50 mg.

Strengthens mucous  
tissue and reduces  
gastric acidity

**ALUMINUM  
HYDROXIDE  
GEL (Dried) U.S.P.**  
100 mg.

Relieves the burning  
and protects  
gastric mucus

**SUPPLIES:** 100 and 1,000 capsules.

Literature and samples  
on request.



*Kremers-Urban  
Pharmaceuticals  
Chemical Division*

*New*

# Terra

Brand of **TERRAMYCIN ELIXIR**

provides: **TERRAMYCIN:**

250 mg. per teaspoonful (5cc.)

**CHERRY-COLOR APPEAL**

**MINT FLAVOR**

*unique*

Terrabon\* presents the only broad-spectrum antibiotic available as an elixir and in such potency. Since 1 teaspoonful (5cc.) is equal in Terramycin content to one 250 mg. capsule, *therapy with the elixir does not require new and unwieldy dosage schedules.*

*The preferred dosage form for*

Pediatric Patients

Geriatric Patients

"Sensitive" Patients who experience difficulty in taking the customary forms of oral antibiotic medication.

\*Trade Mark

**Pfizer**

# bon

*The palatability and  
pharmaceutical elegance of  
Terrabon add appeal to the  
broad antimicrobial activity,  
wide clinical effectiveness  
and high toleration of*

## *Terramycin—newest of the broad-spectrum antibiotics*

*Terrabon suggested for: acute pneumococcal infections, including lobar pneumonia, bacteremia; acute streptococcal infections, including erysipelas, septic sore throat, tonsillitis; acute staphylococcal infections; bacillary infections, including anthrax; urinary tract infections due to E. coli, A. aerogenes, Staphylococcus albus or aureus, and other Terramycin-sensitive organisms; brucellosis (abortus, melitensis, suis); hemophilus infections; acute gonococcal infections; lymphogranuloma venereum; granuloma inguinale; primary atypical pneumonia; typhus (murine, epidemic, scrub); rickettsialpox.*

*Dosage:* One teaspoonful (5 cc.) of Terrabon is equal in Terramycin content to one 250 mg. capsule. In adults and older children, 2 to 3 Gm. daily in divided doses q. 4 or 6 h. is suggested. In children under 20 Kg. body weight, suggested dosage is 100 mg./Kg. body weight per day.

**SUPPLIED** Terrabon is available as a combination package consisting of:

- 1-A vial containing 1.5 Gm. of Terramycin as Terramycin Hydrochloride.
- 2-A bottle containing 1 fluid ounce of specially buffered and flavored diluent.

**ALSO AVAILABLE:** Crystalline Terramycin Hydrochloride in 3 convenient capsule sizes:

250 mg., bottles of 16 and 100;  
100 mg., bottles of 25;  
50 mg., bottles of 25.

*Antibiotic Division*


**CHAS. PFIZER & CO., INC., Brooklyn, 6, N. Y.**

# 3

reasons

why

•  
•



relief  
is  
prompt  
in  
diarrhea



- **KAOLIN**—recognized for centuries as a most effective adsorbent and demulcent;
- **AUGMENTED BY A SPECIAL ALUMINA GEL**—ideal as a base and providing outstanding added protective, soothing, and adsorptive properties;
- **WITH PECTIN**—contributing bland bulk in addition to its soothing, local colloid action. Pleasant-tasting.

## **KAOMAGMA<sup>®</sup>** **WITH PECTIN**

provides the benefits of three clinically established coating agents; quickly restores the patient's comfort.

Bottles of 12 fl. oz.

*Wyeth* Incorporated, Philadelphia 3, Pa.



# MODERN MEDICINE

## *Special Article*

### Physical Medicine and Rehabilitation in Poliomyelitis

ROBERT L. BENNETT, M.D.\*

*Georgia Warm Springs Foundation, Warm Springs, Ga.*

*Prepared for Modern Medicine*

THE general practitioner is frequently called on to care for the patient with the after effects of acute anterior poliomyelitis. Above all else, the physician who accepts this responsibility must realize that the end results depend not only on the site and extent of damage to the central nervous system by the virus during the acute stage, but on what he does with what is left intact and recoverable during all the remaining years of the patient's life.

Even when only slight weakness remains after the acute stage, the problem of regaining effective use of skeletal musculature and of maintaining good skeletal alignment throughout the patient's lifetime is difficult. The only solution lies in a sound knowledge of normal bodily mechanics, long experience in the effects of faulty bodily mechanics on muscles, bones, and joints, and appreciation of the psychic trauma which follows the patient's recognition of his physical limitation and deformities.

These basic requirements, coupled with a deep sense of responsibility to see the patient through the many years of recovery and adaption, are essential to adequate care. Unless these requisites can be met, the physician must not take

\* Director of Physical Medicine, Georgia Warm Springs Foundation.



over a problem so difficult and exacting as that presented by the after effects of poliomyelitis. Some points of basic importance are outlined briefly in the following pages.

*The child with poliomyelitis presents far more difficult problems than the adult with comparable involvement.*

In the adult, the basic problems are restoration of muscle strength and function, followed by resumption of activity within the limits imposed by the residual disability and by the future security of skeletal structures, and the compromise and adjustment necessary to return to a normal environment. Obviously, the problem varies with the extent of involvement but in all phases is clearly outlined.

The child has all these difficulties and, in addition, that of skeletal growth. In the presence of muscle weakness, the effect on bone growth of changes in the child's height, weight, and scope of activity is the most difficult problem in the care of poliomyelitis.

The general practitioner must recognize this fact and guide the patient and his family over the long years that are required for optimum results. Even when involvement is apparently slight, the growing child is constantly faced with the possibility of skeletal deformity. Practically all these deformities can be minimized if recognized early and wisely treated.

*Poliomyelitis is capable of producing changes not only in muscle strength and coordination, but also in supportive tendinous, ligamentous, and fascial tissues.* These changes usually result in loss of joint mobility through contractures, but may cause loss of stability through relaxation.


Tight tendinous structures may produce marked limitation of muscular strength and faulty bodily mechanics. Likewise, relaxation of these structures can cause deformity and incomplete return of muscular function. The physician must be able to recognize not only strength and weakness but also tightness and relaxation so that he can balance one against the other to encourage return of functional strength in intact muscle tissue and, at the same time, not endanger stability by stretching structures that have little chance of recovering necessary muscular support.

*A fairly constant relationship exists between the time that has elapsed since the acute onset and the extent of recovery*



of muscle function that is possible. The gross expectancy in any period of time is indicated in the chart.

Proper appreciation of this time and recovery relationship makes it possible to continue treatment as long as there is significant hope of further recovery and to avoid wasting time and money in the vain hope of recovery beyond a reasonable expectancy. For example, if a muscle group is rated "poor minus" or 10% of normal strength at the end of six months of adequate treatment, this score represents approxi-



mately 75% of the expected return. The additional 25% recovery possible through treatment over an indefinite period of time would not appreciably change the functional rating of the muscle and would only bring up the percentage to 12½% of normal. This gain would be insufficient to justify intensive treatment beyond the period when no statistical hope of real function remains.

On the other hand, if the muscle group is rated "fair" or 50% of normal at the end of six months, a period of treatment for a year or even longer would be well worth while if, at the end of that time, the additional 25% of recovery expected would bring the muscle up to a "fair plus" or even "good minus" grading.

The chart refers to the recovery of individual muscle

power through muscle reeducation and does not reflect the possibilities of gain in independence through specific functional training. The chart has its greatest value in the analysis of muscle grades under the 50% or functional level.

*The period of recovery is divided into two fairly distinct phases.* The first phase consists of individual muscle reeducation based on the assumption that, with time and good treatment, sufficient strength and coordination will return to permit normal activities. During this phase, only normal patterns of muscle action are attempted and no substitute or incoordinate patterns are permitted.

When time and good treatment indicate that recovery will not be complete, some compromise in normal activities may be made. Functional patterns of substitution may not only be permitted but actually encouraged as the patient is gradually taught to use what muscle strength has been recovered to specific functional effect. This is the second phase, or the phase of functional training. Any attempt to initiate this phase before muscle reeducation has brought as much recovery as is possible in the individual muscle or muscle group can only result in incomplete attainment of the maximum functional end result.

*Persistent faulty alignment of bodily segments during rest or activity is the major cause of musculoskeletal deformities.*

The alignment of bone is largely dependent upon the stresses placed on it. Abnormal stresses such as faulty muscle balance and usage, contractures of muscular, ligamentous, and fascial tissues, lack of skeletal support or improper support, overweight, overactivity, and so on, must be recognized and controlled early if deformities are to be prevented or kept at a minimum.

Anyone with but little medical experience and training can recognize far advanced deformities. However, once this point is reached, little can be done beyond radical orthopedic surgery. The physician who cares for the patient with poliomyelitis must, therefore, be prepared to recognize these factors before they bring about irreversible changes and to follow through with a program of care designed to control the stresses before the deformity progresses so far that it cannot be kept at a minimum by conservative care and properly timed surgery.

*Apparatus is frequently a necessary part of convalescent as well as of chronic care.* Splints, corsets, crutches, braces, and so forth, may be essential, not only in the support of weakened bodily segments and in the prevention or correction of musculoskeletal deformities but also in muscle re-education.

The physician must, therefore, be able to recognize the need for such apparatus and to write an accurate prescription for the brace-maker. Much of the present feeling against the use of apparatus in the convalescent stage is based on an inadequate knowledge of bracing methods by physicians in charge of treatment and their inability to obtain well-fitting splints and braces from near-by manufacturers. For the severely involved patient, the physician's failure to appreciate proper apparatus and to prescribe accurately and obtain such apparatus may well mean the difference between independence and complete helplessness.

*Most patients, particularly children, require a long period of treatment at home.* This treatment usually consists of some combination of specific strengthening and stretching routines, limited physical activity, and proper use of necessary apparatus. Unless the patient or his parents have the intelligence and responsibility to carry out basic routines, the value of professional care will be lost.

Obviously, the physician in charge must be able not only to prescribe such care and realize the importance of its being carried out over long periods of time but he must also be able to determine that the therapy is being carried out. If it is evident that the child can never receive adequate treatment at home and his disability is such that treatment is necessary, some radical compromise must be made in bracing and possibly early surgery to obtain some part of the expected result. If the parents and the patient are irresponsible there is little advisability of long hospital care when its value will only be dissipated by negligent and haphazard treatment at home.

*The physician must recognize the value of properly timed orthopedic surgery in the overall program.* Surgical procedures designed to release persistent contractures or remove the deforming pull of muscle imbalance may be necessary in the convalescent stage not only to restore bodily alignment and

avert severely handicapping structural changes but also to allow the most effective use of muscle reeducation and functional training. Obviously, then, the physician must know the indications for these procedures and must not temporize with conservative routines when more radical procedures are necessary.

*All patients with longstanding and apparently hopeless residual handicap deserve periodic evaluation.* The field of physical medicine and rehabilitation is constantly developing routines and apparatus that may bring about startling functional improvement in patients formerly considered beyond further help. It is difficult to imagine a condition so chronic that the patient cannot be helped by thoughtful evaluation and specific treatment.

## Adductor Longus Syndrome

JANET TRAVELL, M.D.\*

SPASM and tenderness of the adductor longus muscle may cause persistent pain in the groin and leg. Relief may be obtained by anesthesia of the most sensitive areas, finds Janet Travell, M.D., of Cornell University, New York City.

In some cases the syndrome is a physiologic disorder of myofascial tissues, in others, a reflex cycle set up by remote lesions. Pain referred from the upper half of the muscle is felt deep in the groin. Sensations from the lower part radiate over the anteromedial aspect of thigh and knee, and at times over the pretibial region to the ankle.

Ethyl chloride spray is applied lightly over the muscle and groin in rhythmic upward sweeps, with avoidance of extreme chilling. During treatment the muscle is gently stretched. If tension does not relax immediately, the trigger zone should be infiltrated with 0.5% procaine hydrochloride in physiologic saline solution.

Symptoms due to direct muscle injury are permanently abolished by anesthesia. If spasm arises from a distant lesion, such as carcinoma of the lumbar spine or osteoarthritis of the hip, remissions are temporary, as a rule. Some patients apply the spray at home with good effect.

\* The adductor longus syndrome: a cause of groin pain; its treatment by local block of trigger areas (procaine infiltration and ethyl chloride spray). *Bull. New York Acad. Med.* 26:284-285, 1950.

# Treatment of Venous Thrombosis

F. RICHARD PIERCE, M.D., AND TRENTO J. DOMENICI, M.D.\*

*Henry Heywood Memorial Hospital, Gardner, Mass.*

**M**ODERN management of venous thrombosis consists of anticoagulants, surgical interruption, or both (see table).

Prevention of pulmonary embolism lies in the early recognition of a venous thrombus. Phlebothrombosis, which has a dangerous propensity for embolization, must be differentiated from thrombophlebitis, with the more adherent clot.

Although both conditions may co-exist, thrombophlebitis is an acute inflammatory process with heat, swelling, tenderness, and a fixed clot. Phlebothrombosis is noninflammatory, produces slight fever, tachycardia, tachypnea, minimal calf tenderness, and ankle edema and is often suspected only after pulmonary infarction has occurred.

In the experience of F. Richard Pierce, M.D., and Trento J. Domenici, M.D., early ambulation of the postoperative patient has not appreciably decreased the incidence of venous thrombosis. The condition, furthermore, is commoner in medical than in postoperative patients.

The use of anticoagulants is directed against the basic lesion throughout the vascular tree, rather than only the venous system of the legs. This method also lessens the possibility of postphlebitis or post-thrombotic sequelae.

Numerous conditions, however, mitigate against the use of anticoagulant drugs, while no contraindication to venous interruption is recognized. Dicumarol, one of the most common anticoagulants used, requires facilities for daily determination of prothrombin time.

By whatever method employed, an effective level exists when the patient's prothrombin time is twice that of the control. For immediate anticoagulant effect, heparin is given intravenously in a continuous 250- to 500-cc. saline infusion with a total dose equivalent to 50 to 100 mg. of the drug every four hours. After the prothrombin time is determined, dicumarol is started at an average dose of 300 mg. the first day, 200 the second, and 100 the third. After this, heparin can usually be discontinued and prothrombin kept between 45 and 60% by dicumarol.

The antidote for excessive heparinization is intravenous 1% protamine sulfate in 50- to 100-mg. doses. Excessive dicumarol effect is best combatted by vitamin K in doses of 68 mg. intravenously every four hours and fresh whole blood transfusions. Anticoagulant therapy should be maintained as long as the patient's symptoms suggest continued activity of the basic process.

Dicumarol used ante partum may

\* Problems and practices in a community hospital: treatment of venous thrombosis. *New England J. Med.* 242:395-402, 1950.

## MEDICINE

produce hemorrhage in the newborn but is beneficial in treatment and prophylaxis of postpartum venous thrombosis.

Venous interruption is the treatment of thrombosis when insufficient laboratory facilities preclude the use of dicumarol.

An adequate femoral vein interruption implies opening and exploring the vein. The danger of loosening a fatal embolism is slight, and removal of the clot protects against recanalization or future embolization.

The operation is performed bi-

laterally, since the apparently unaffected leg often harbors the thrombus. The superficial femoral vein is interrupted unless the clot extends into the common femoral vein, whereupon the latter vessel is explored and the clot aspirated. Surgical obstruction of the common femoral vein or of the inferior vena cava is to be avoided if possible, since impaired collateral circulation and serious disabling sequelae may result. Sympathetic block, Priscol, and papaverine may be used to relieve reflex vasospasm.

INDICATIONS FOR PROCEDURES WITH VENOUS THROMBOSIS

	ANTICOAGULANT TREATMENT	VENOUS INTER- RUPTION
<b>PROPHYLACTIC INDICATIONS*</b>		
<i>Medical</i>		
Myocardial infarction	Indicated	Not used
Prolonged bed rest with history of venous thrombosis	Indicated	Not used
<i>Surgical</i>		
Elderly patients with:		
Major amputations or severe trauma to lower extremities	Not used	Indicated
Extensive gastrointestinal surgery	Not used	Indicated
Stage operations	Not used	Indicated
<b>THERAPEUTIC INDICATIONS*</b>		
<i>Medical</i>		
Prolonged enforced bed rest with development of signs or complications of venous thrombosis (congestive heart failure, pneumonia, prolonged infections)	Indicated	Not used
<i>Obstetric</i>		
Venous thrombosis or its complications in:		
Antepartum cases	Contraindicated	Indicated
Postpartum cases	Indicated	
<i>Surgical</i>		
Phlebothrombosis with slight to moderate or no pulmonary involvement	Indicated	Not used
Phlebothrombosis with massive pulmonary infarction	Indicated†	Indicated†
Thrombophlebitis (to guard against phlebothrombosis in opposite leg)	Indicated	Not used
Venous thrombosis with:		
Impaired liver or kidney function	Contraindicated	Indicated
Blood dyscrasia	Contraindicated	Indicated
Surgery in which hemorrhage or oozing may be postoperative factor	Contraindicated	Indicated
Active bleeding from any source (peptic ulcer, uterine tumor)	Contraindicated	Indicated
Failure of response to anticoagulants	Contraindicated	Indicated
Neurologic disease or injury to central nervous system	Contraindicated	Indicated
Prolonged use of anticoagulants, making them expensive or impracticable	Contraindicated	Indicated
Prolonged use of acetylsalicylic acid	Contraindicated	Indicated
Patients with pulmonary infarction		
When heparin is not immediately available	Contraindicated	Indicated
With repeated emboli in spite of anticoagulants		Indicated

\* If laboratory facilities are inadequate, patients with venous thrombosis or its complications should have venous interruption.

† Veins interrupted as emergency measure with simultaneous administration of anticoagulants.



# Current Histamine Therapy

LESTER S. BLUMENTHAL, M.D.\*

George Washington University, Washington, D. C.

**I**N proper dosage, histamine is a safe drug and is of definite benefit in Ménière's syndrome, histaminic cephalalgia, selected cases of migraine headache, nonspecific photophobia, and acute and chronic urticaria.

Lester S. Blumenthal, M.D., has administered histamine with no untoward effects to patients with peptic ulcer, bronchial asthma, or other conditions for which the drug has ordinarily been considered hazardous.

The antihistaminic drugs, by blocking specific histamine effects, actively interfere with and modify typical allergic and anaphylactic processes. In the allergic reaction, histamine is probably not the primary factor involved, but is the chemical released secondary to some disruptive process within the cell. Therapeutic applications of the drug are based on the assumption that some fundamental sensitivity exists that can be neutralized by desensitization.

**Histaminic cephalalgia**—Horton's headache seems related to histamine since [1] a subcutaneous injection of histamine base usually precipitates an attack within thirty minutes and [2] attacks generally cease when gradually increasing subcutaneous doses of histamine are given twice daily for ten to thirty days.

Histamine may be given subcutane-

ously or intravenously. About a fifth of patients require no further treatment after a course of intravenous therapy. For the others, however, desensitization must be completed subcutaneously.

For subcutaneous dosage histamine diphosphate, 0.275 mg., equivalent to 0.1 mg. of histamine base, is available in 1-cc. ampules. At the first injection, 0.25 cc. is given, and subsequent doses are each increased by 0.05 cc. until the patient is receiving 1 cc. as the sixteenth injection.

If subjective or objective effects are noted, the next dose is cut in half. Eventually the amount is found that invariably produces symptoms. This represents the tolerance dose. More than 1 cc. should not be given to any patient.

The foregoing desensitization regimen usually improves or alleviates symptoms but, for prophylactic purposes, the frequency of injections should be gradually decreased. Eventually further treatment is unnecessary or the patient may require a tolerance dose at regular intervals, such as once every one or two weeks. If symptoms recur after a year or two, the original regime may be repeated.

For intravenous dosage a specially prepared solution is used, each 100 cc. of which contains 0.55 mg. of

\* Current histamine therapy. M. Ann. District of Columbia 19:191-198, 1950.

## MEDICINE

histamine diphosphate and 0.9 gm. of sodium chloride.

Because of the potential effect on gastric acidity, treatments are administered immediately after eating and antacid tablets are chewed continuously during the injection. The procedure lasts an hour and may be done at the office.

The solution must not be run in too rapidly. Each injection is started at 5 to 10 drops per minute and the rate is gradually increased until a slightly flushed or full sensation in the head is barely perceptible. After several treatments, 40 to 60 drops per minute is satisfactorily tolerated. The injections are given daily for four days, then every other day.

As soon as definite improvement is noted, the subcutaneous route may be substituted, and the patient administers the drug to himself at home.

For 20 patients with histaminic cephalalgia, the intravenous route was employed and in most cases great relief was obtained within several days.

*Ménière's disease*—Intravenous histamine is effective for acute attacks of Ménière's disease.

To avoid introduction of the sodium ion and because potassium chloride often causes severe burning sensations along the vein, a 5% solution of dextrose in distilled water is used as the vehicle; 1 ampule of histamine diphosphate, 2.75 mg. per cubic centimeter, is diluted in 500 cc. of the dextrose solution. The speed of injection is gradually increased until the skin flushes, usually at the rate of 50 to 60 drops per minute.

A one-hour treatment is given daily until headaches are relieved, ordinarily in three days, after which subcutaneous injections are employed prophylactically.

Therapy of the acute attack is more satisfactory than the prevention of recurrences.

*Migraine headache*—The principal use of histamine in treatment of migraine is to prevent or reduce the incidence of headaches.

Cases in which benefit may be expected are those with a unilateral, periodic, throbbing, vascular type of headache, associated with two or more of the following: [1] gastrointestinal dysfunction, [2] prodromes of scotomas or other cerebral disturbance, [3] hemicrania among at least two other members of the family, [4] duration of the usual attack for thirty-six hours or more if no medication is given, and [5] complete freedom from pain at intervals.

With one-hour intravenous injections given every other day for one to two months, followed by intravenous injections at intervals of seven to ten days or subcutaneous therapy, approximately 300 patients had 75% improvement.

*Photophobia*—Since an abnormal intolerance to light is often associated with migraine, 3 patients with photophobia were treated with intravenous and subcutaneous histamine. Results were excellent.

*Posttraumatic syndrome*—Vasodilating headache is a frequent complication of the posttraumatic cerebral syndrome. Daily intravenous histamine for about two weeks brought definite early relief of symptoms to 3 of 5 patients and more gradual



but progressive improvement in the other 2.

*Acute and chronic urticaria*—For 5 patients with acute postmedication urticaria who had not benefited from previous treatment with epinephrine, calcium, or antihistamine, complete recovery within twenty-four hours of the last injection was achieved with histamine. Intravenous administration of 30 mg. of Benadryl was followed by rapid injection of 500 cc. of dextrose solution containing 1 mg. of histamine base daily for three days.

Of 9 patients with chronic urticaria treated by combined intravenous and subcutaneous histamine, all showed definite improvement although results were not as striking or as rapid as in the acute cases.

*Miscellaneous conditions*—Daily in-

travenous histamine injections were given to 4 patients with sudden nerve deafness presumably resulting from edema of the cochlear portion of the labyrinth. Best results were attained in the most recent cases.

Considerable benefit for symptoms of obliterative peripheral vascular disease followed biweekly infusions of histamine into the femoral artery.

Coagulation time is shortened as a result of increased platelet activity in patients receiving intravenous injections of histamine. For 5 patients with active hemophilia given intravenous histamine, cessation of bleeding, an increase in well-being, and a decrease in coagulation time were noted.

Intravenous histamine therapy appears to be of value in the treatment of temporal arteritis.

**L**ABIAL DIPHTHERIA may develop alone or with lesions in other sites. During a widespread epidemic in a Japanese prison camp on Singapore Island, 46 British prisoners had typical membranes on the lips, particularly at corners of the mouth. Riboflavin deficiency with cheilosis and fissure was the predisposing factor, says George S. Riddell, M.D., of the Aberdeenshire County Council, England. Simultaneous nasal, faucial, or cutaneous infection appeared in only 22 instances.

*Brit. M. J.* 4637:818-819, 1950.

**A**EROBIC FECAL BACTERIA appear to be reduced only temporarily, and anaerobic organisms not at all, by sulfonamide therapy. In 1 case of chronic ulcerative colitis reported by Homer C. Marshall, Jr., M.D., Joseph B. Kirsner, M.D., and Walter L. Palmer, M.D., of the University of Chicago, the flora changed from predominantly gram negative to predominantly gram positive while ulceration of the bowel persisted, and returned to gram negative by the time the condition had healed. Only 2 of 5 patients showed clinical improvement during treatment.

*Gastroenterology* 14:418-424, 1950.

**D**IAGNOSIS OF ADDISON'S DISEASE may be confirmed by adding a few drops of Millon's reagent to serial dilutions of urine. If the color changes to pink, Addison's disease is indicated. When reactions are equivocal, the test is repeated with urine obtained after ingestion of 4 gm. of tyrosine. C. Ferrero, M.D., of Geneva University, Switzerland, found reactions positive in each of 13 cases of Addison's disease and in 9 of 12 cases of hypopituitarism with secondary adrenal insufficiency. In 13 other endocrine disorders, the reaction was positive only in hypothyroidism. A negative reaction is obtained when desoxycorticosterone acetate or adrenal extract is being given, and return of a positive reaction during treatment indicates insufficient medication.

*Schweiz. med. Wchnschr.* 80:179, 1950.

**L**IVER EXTRACT INJECTIONS are sometimes discontinued because of intolerance. However, desensitization may be achieved by a series of progressively larger injections, administered by the patient or a member of his family at home. Sensitivity reaction usually consists of flushing, nausea, hives, or other allergic symptoms. Oral liver therapy may be tried but frequently is incapable of preventing pernicious anemia relapse. Changing the animal source of the liver extract usually is of no benefit. Beginning with 0.05 cc. liver extract, the dose is increased daily by 0.05 cc. until 0.6 or 0.8 cc. can be given at one time. Jerome E. Cook, M.D., of Washington University, St. Louis, then instructs the patient to take a small injection of liver extract every two or three days; the exact dosage depends upon the patient's hematologic needs. Once the patient has become desensitized, liver extract injections must be given at frequent intervals, such as every other day, or reactions may recur.

*Ann. Int. Med.* 32:506-509, 1950.

**A**BSOLUTE LYMPHOCYTOPENIA within forty-eight hours of onset of the disease supports a diagnosis of acute pancreatic necrosis. A later rise in the number of lymphocytes allows a good prognosis. A diagnosis of acute necrosis was made in 38 cases by K. Herfort, M.D., and V. Letosnik, M.D., of Charles University, Prague. All the patients had lymphocytopenia; the number of lymphocytes in the peripheral blood decreased to from 2 to 10% of the total white count, which remained normal or increased only slightly. In 31 cases, diagnosis was confirmed by operation or necropsy. The lymphocyte values of patients showing improvement returned to normal levels.

*Casop. lek. cesk.* 88:1006-1009, 1949.

## Commissurotomy for Mitral Stenosis

ROBERT P. GLOVER, M.D., THOMAS J. E. O'NEILL, M.D.,  
AND CHARLES P. BAILEY, M.D.\*

*University of Pennsylvania, and Hahnemann Medical College, Philadelphia*

CONSIDERABLE function may be restored to a stenotic mitral valve by separating the leaflets along the fused border.

The surgeon's forefinger is inserted through the auricular appendage, and a hooked knife is slipped down the finger and manipulated by the sense of touch. For unusually hardened tissue, Robert P. Glover, M.D., Thomas J. E. O'Neill, M.D., and Charles P. Bailey, M.D., use a specially designed punch with trocar and cannula.


Pressure in the left auricle is immediately reduced without increase of regurgitation. Invalids are often able to give up digitalis and return to former occupations. Of 30 persons undergoing the procedure, 21 were greatly benefited and 3 improved subjectively. Half of the 6 deaths occurred before surgical technic was perfected.

Commissurotomy is done only in carefully chosen cases. Recovery is most likely if the mitral defect produces extreme fatigability, exertional dyspnea, and definite pulmonary

hypertension. Rheumatic disease should be quiescent, the valve primarily stenotic, with little incompetence, and sinus rhythm normal. The left ventricle should not be appreciably enlarged.

Prospects are less encouraging for patients with recurrent blood stains in sputum, arterial emboli, or auricular fibrillation without heart failure.

Operation is contraindicated by rheumatic infection, subacute bac-



\* Commissurotomy for mitral stenosis. *Circulation* 1:329-342, 1950.

## SURGERY

terial endocarditis, mitral regurgitation, aortic valve deformity, or heart failure beyond medical control.

The heart is approached through the left anterior chest wall with the patient in dorsal recumbent position. The pericardium is incised longitudinally on the left lateral aspect,  $\frac{1}{2}$  in. anterior to the phrenic nerve.

The huge auricular appendage now protruding from the pericardial sac is encircled at the base with a purse-string suture of heavy braided silk. A Satinsky clamp is closed over the base, and a generous portion of the tip is amputated.

The operator wears two gloves on the right hand. The hooked blade is inserted at the base of the index finger through a slit in the outer glove and projected at the tip.

The finger is slipped into the left auricle as the clamp is released and as the purse-string suture is pulled taut. The valve is found with the finger tip, and size of the opening and location of the commissures are noted.

The knife is then thrust through the valvular orifice, and the antero-lateral commissure is divided with a backward stroke of the hook. The finger again palpates and gently di-

lates the opening, and if the cut does not extend well into healthy tissue, the knife stroke is then repeated.

When the valve is rigid and fixed the incision may be widened. The hand is pronated to divide the medial commissure.

As the finger and knife are withdrawn from the auricle, the purse-string suture is drawn tight, so that only a few cubic centimeters of blood is lost. The suture is tied and the cut edge of the appendage oversewn.

The backward-cutting punch designed for very hard valves has a narrow blade and long bite. The incision can be extended through diseased tissue into the flexible membranous valve as with the hook, but has not the advantage of finger guidance.

Postoperative pain is reduced by opiates for three or four days, and walking is allowed in two to five days.

As shown by cardiac catheterization, reopening the mitral valve at once lowers pressure in the chambers. The systolic value in the left auricle may fall from 45.1 to 15.5 mm. of mercury and diastolic pressure from 30.3 to 2.6 mm.

**R**ESPIRATORY BORBORYGMUS may be present with hiatus hernia and possibly with other conditions in which the stomach is partly in the thorax. The sounds are usually found approximately at the level of the diaphragm. Although gurgling sounds in this area are not uncommon, the borborygmi described by Henry J. Marriott, M.D., of University of Maryland, Baltimore, in hiatus hernia were rhythmic and synchronized with respiration. The sounds are intensified when the patient is in a sitting position.

*Bull. School Med. Univ. Maryland 35:13-15, 1950.*

## Thymectomy for Myasthenia Gravis

HENRY R. VIETS, M.D.\*

*Harvard University, Boston*

RESULTS of thymectomy in the last nine years justify use of the operation for treatment in about half the cases of myasthenia gravis. If patients are carefully selected and adequately maintained with neostigmine during the procedure, operative mortality should be negligible.

Thymectomy should be performed only when the patient is steadily and fully maintained and well balanced by neostigmine because of the hazard of remissions and relapses. Operations should never be done hastily.

Thymectomy is not the best therapy for patients with few symptoms, those well benefited by neostigmine regardless of the dosage, most persons over forty years of age, or for patients with wide swings of remission and relapse. Less than half of 300 patients seen since 1935 by Henry R. Viets, M.D., have been considered proper candidates for thymectomy.

Since 1941, of 36 myasthenic patients subjected to thymectomy, 7 had thymomas and 29 had various degrees of involution of the thymus with germinal centers in thymic tissue.

Results of the 25 survivals in cases without thymoma were classified as excellent to good, 10; fair to good, 4; poor to fair, 2; poor, 2; and too soon to estimate, 7.

The conditions of 2 of the 3 living thymoma patients are excellent, that of the third is fair; 2 of the 4 nonsurvivors were operative deaths, presumably from improper handling at operation. Neither patient had the advantage of a properly adjusted amount of intravenous neostigmine.

The appearance on a radiogram of a mass in the anterior mediastinum in a myasthenic patient indicates probable existence of a thymoma. Operation may or may not be advised, the decision being based on a careful study of the entire condition. Some thymomas become invasive but are usually considered benign neoplasms. Implantations may occur postoperatively.

Dosage should be fixed for a week or more before the operation. Each 15-mg. oral tablet of neostigmine bromide is the equivalent, in effect, of 0.5 mg. of neostigmine methylsulfate given intravenously. Thus, a patient taking 12 tablets in twenty-four hours requires 2 mg. of neostigmine methylsulfate intravenously in an equivalent eight hours. This may be added to 1,500 cc. of dextrose or saline and given intravenously just before, during, and after the operation.

Diminution of the severity of relapses and prolongation of duration of remissions following thymectomy

\* Thymectomy in myasthenia gravis. *Brit. M. J.* 4646:139-147, 1950.

## SURGERY

are often favorable effects of the operation. A patient's postoperative status is more stable than during the preoperative period.

Age, sex, duration of disease, length of previous treatment with

neostigmine, roentgen-ray treatment, and condition of the thymus at operation are not correlated with the degree of improvement after thymectomy. Symptoms may improve immediately or after several months.

## Improved Biliary T-Tube

R. RUSSELL BEST, M.D.\*

A RIGHT-ANGLE tube, though useful for routine postoperative or delayed cholangiographic studies, is not suitable for flushing out residual stones, finds R. Russell Best, M.D., of the University of Nebraska, Omaha. Solutions injected through the long arm often pass toward the liver and not into the lower end of the duct, where debris generally collects.

A tube with a wide angle and second lumen directs fluid to the desired site (see illustration). The double lumen permits escape of gas and fragmented debris and allows two-way irrigation.

The long arm of the modified tube forms an angle of  $135^{\circ}$  with the arm that enters the distal part of the common duct. The device is more easily introduced and occupies a better anatomic position than the usual model.

The double lumen of the tube runs down the long arm along the wide angle to the distal end of the transverse arm. The smaller of the two channels projects from the side of the long arm and fits an ordinary syringe tip.

When the T-tube is inserted, a heavy silk suture is passed through one of four rubber tabs, according to abdominal depth, and through the skin. The tube is thus firmly fastened, yet the lumen is not punctured. Adhesive tape may be added for security.

Antispasmodics and solutions to wash out or dissolve stones are introduced into the smaller opening by a syringe with blunt needle. Contrast media are injected through the large lumen, while the second passage is plugged or compressed.

\* Cholelithotomy—advantages of a modified T tube. *Surg., Gynec. & Obst.* 90:295-297, 1950.



## Ulcerative Colitis

J. M. RICE-OXLEY, B.M., AND SIDNEY TRUELOVE, M.D.\*

*Oxford University, England*

IN spite of great advances in other medical fields, ulcerative colitis remains dangerous and disabling. Treatment and prognosis vary with the stage of the disease.

Mortality is 22% in the first year and then declines, although 10% of cases may prove fatal in the next five years. During the fulminating state, even when medical measures are apparently failing, emergency ileostomy seems to do no good and may precipitate death, observe J. M. Rice-Oxley, B.M., and Sidney Truelove, M.D.

If the patient survives the first year of the disease, colitis usually recurs, sometimes after a long remission. At least half the patients are more or less incapacitated and, in such instances, ileostomy may be justified even in uncomplicated cases.

Estimates of probable outcome and of the value of specific remedies for ulcerative colitis are often unreliable because statistics are based on a heterogeneous group of patients admitted to the hospital in all phases of involvement. Moreover, investigators frequently fail to distinguish between acute and chronic disease.

To obtain a representative sample, 72 patients with ulcerative colitis were observed for one to eleven years, starting with the first attack. All subjects had been ill less than two years when first admitted to a general hospital.

Medical measures consist chiefly of rest in bed; replacement of blood, electrolytes, and protein by intravenous infusions; low-residue diets high in calories and proteins; and vitamin supplements.

Chemotherapy and antibiotics apparently do not change the course of ulcerative colitis. However, penicillin and sulfonamides have some value if used before and after operations and for prevention or treatment of pyogenic complications.

After the acute stage, the disease tends to become intermittent, with remissions of five years or more, but in some cases symptoms continue indefinitely.

A number of patients managed to lead practically normal lives regardless of diarrhea. About half are moderately incapacitated, with frequent admissions to the hospital, and a small proportion are totally disabled. Percentages in each group are about the same whether colitis is continuous or interrupted.

Since medical treatment is unsatisfactory and the relapse rate high, inconveniences of ileostomy should be carefully weighed against the discomforts and restrictions of chronic disease. Though generally considered only for a stricture, fecal fistula, or other complication, operation may be advisable in cases of several years' standing.

\* Ulcerative colitis: course and prognosis. *Lancet* 258:665-666, 1950.

## **more physicians are satisfied**

The development of the new improved Biolac supplies a long-sought need in infant nutrition. To accomplish this, Borden scientists surveyed our present nutritional knowledge. They then tested more than 500 formulations. Having decided on the formula that would best supply the normal infant's nutritional requirements in their most assimilable form, a modern plant was constructed in 1949 so that the new formula could also benefit from the most up-to-date techniques and control in processing equipment. A Biolac formula that is both new and improved is thus made available.


Biolac is intended for prescription by every physician with infants among his patients. It satisfies the physician's demand for a complete food to which only vitamin C need be added. That means it is simplicity itself to prepare and provides the maximum in formula safety for the infant.

And yet, for all these advantages, Biolac costs no more.



**The Borden Company**





For up-to-date, complete  
infant nutrition, prescribe *new improved*

**Biolac**<sup>®</sup> a development of

The Prescription Products Division

The Borden Company

*Ingredients:*

nonfat dry milk solids, dextrins-maltose-  
dextrose, lactose, coconut oil, destearinated beef fat,  
lecithin, sodium alginate, disodium phosphate, ferric  
citrate, vitamin B<sub>1</sub>, concentrate of vitamins A and D from fish liver  
oils, and water. Homogenized and sterilized.

Dilution: one fluid ounce to one and a half ounces of boiled  
water for each pound of body weight.

Biolac is available at drugstores in 13 fluid ounce tins.

Prescription Products Division, 350 Madison Avenue, New York 17

## Elastic Hand Splints

STERLING BUNNELL, M.D., AND LOT D. HOWARD, JR., M.D.\*

San Francisco

**A**FTER injury of the hand, stiffened joints can be coaxed into a new position by elastic or spring splints.

Several types are employed for exercise or mobilization, to restore a position of function, or, in case of paralysis, to complete the muscle balance. Sterling Bunnell, M.D., and Lot D. Howard, Jr., M.D., have recently invented new models and supplied attachments for splints already in use.

The object of the first adaptation is to change the hand from a non-functional status, with wrist flexed and fingers extended, to the grasping attitude, with wrist dorsiflexed and fingers bent. The basic knuckle-bender splint is employed with three new attachments.

Figure *a* shows the hand in the position of nonfunction, with the splint and two supplements in place. In Figure *b* the hand has been drawn into a position of function.

The first attachment is a flat padded segment hinged to the distal part of the splint to flex the middle finger joints. Both old and new sections are powered by rubber bands.

The second addition is an Oppenheimer type of splint, which slowly forces the wrist into dorsiflexion by the gentle elastic pressure of a spring. The two attachments are fastened to the original

splint by wire rods thrust into metal sleeves.

The third device, a spring cock-up splint, can be substituted for the Oppenheimer splint to dorsiflex the wrist. The cock-up splint has a notched hand piece to fit over the crossbar of the knuckle-bender, as illustrated in Figure *c*.

Either the Oppenheimer or the cock-up attachment can be used with the knuckle-bender alone or in combination with a device to flex or straighten the fingers.





A miniature replica of the knuckle-bender splint is worn for a single stiff finger joint, leaving the rest of the hand unencumbered. In Figure *d* the finger wearing the splint is still extended, in Figure *e* the middle joint is being flexed. Elastic tension is maintained by druggist's light-weight rubber bands. The proximal wire extension is bent to accommodate the finger webs.

Still another splint draws out flexion contracture of the finger joints after ischemic fibrosis of hand muscles. The proximal segments alone are extended by a knuckle-bender in reverse. Figure *f* shows the deformed position with the splint in place; in Figure *g* the proximal finger joints have been pulled into extension by three-point elastic pressure.

A base plate is stabilized in the

\* Additional elastic hand splints. *J. Bone & Joint Surg.* 32-A:226-228, 1950.

## APPLICATION OF SPLINTS



palm by a web strap fastened about the wrist. Counterpressure is applied over the proximal joints by a slightly curved plate padded with felt.

Elastic pressure is brought to bear on the volar aspect of the finger segments near the distal ends by a padded base. Extended wire outriggers provide good mechanical ad-

vantage and a convenient support for rubber bands.

A difficult problem is thus solved, inasmuch as severe deformity of the type illustrated usually requires an operation. Splints applying elastic extension to the finger tips merely accent hyperextension of the middle and distal joints.

## Pulp Space Infection of the Finger

HAROLD BOLTON, M.D. AND ASSOCIATES\*

**B**y the time a physician is consulted for fingertip infection, an abscess has almost certainly formed. The cavity should be drained at once through a small incision made directly over the site.

Lateral J incisions and bilateral linear cuts often traverse and infect healthy tissue, heal too fast for adequate drainage, or sever the nerves and produce causalgia.

Harold Bolton, M.D., P. J. Fowler, M.D., and R. P. Jepson, M.D., observed superficial or deep central or lateral abscess in all distal phalanges incised at the Manchester Royal Infirmary, England. In 39 of 50 cases the pulp alone was affected, in the remainder bone was apparently involved.

The site of greatest tension and tenderness is located with a blunt probe, or the abscess is found by a spontaneous sinus, which is always directly over the cavity. Bleeding is prevented by a pneumatic tourniquet.

General anesthesia is used and a small ellipse of cornified epithelium is removed. If a small sinus has formed in the true skin, the opening is excised elliptically for a distance of 4 or 5 mm.

Since the wound is open, no drainage is required. The cavity is filled with penicillin in lactose powder and covered with broad-mesh petroleum jelly gauze and a layer of dry gauze. Dressings are changed two or three times at three-day intervals.

If bone infection is suspected or proved, penicillin is injected intramuscularly in doses of 100,000 units twice daily for ten to fourteen days.

Without osteomyelitis, the wound usually heals in ten or twelve days and even a typist can resume work within three weeks.

\* Natural history and treatment of pulp space infection and osteomyelitis of the terminal phalanx. *J. Bone & Joint Surg.* 31-B:499-504, 1949.

## Frequency of Plural Births

METROPOLITAN LIFE INSURANCE COMPANY\*

New York City

**A**GE and race influence the frequency of multiple births. The rate of such births rises with the increasing age of the mother to a peak between the ages thirty-five to thirty-nine and then falls gradually, finds the Metropolitan Life Insurance Company.

While this curve is the same for both white and nonwhite mothers, the incidence of plural births is

This high proportion of multiple births among the nonwhite races is almost entirely a reflection of the rate among Negroes. Other nonwhite groups have a lower incidence of plural births than the white race.

Statistics on multiple births of more than 4 children are, of course, nonexistent. Only two sets of quintuplets are known to have survived infancy—the Dionne quintuplets,

MULTIPLE BIRTHS IN UNITED STATES, 1934-47

Age and Color of Mother	Number of Confinements	Multiple Births per Million Confinements			
		Total	Twins	Triplets	Quadruplets
Total, all ages	35,985,990	10,997	10,889	106	1.6
Under 20	4,276,546	6,178	6,137	41	†
20-24	11,294,105	8,600	8,531	68	1.1
25-29	9,862,737	11,342	11,240	101	1.2
30-34	6,234,130	14,366	14,207	157	2.2
35-39	3,278,517	17,247	17,015	228	4.6
40-44	956,597	14,085	13,916	166	3.1
45 and over	83,358	8,757	8,661	84	†
White, all ages	31,624,779	10,688	10,589	98	1.1
Nonwhite, all ages	4,361,211	13,236	13,062	169	5.0

† Cases too few to warrant computation.

greater at every age for nonwhite mothers. The statistical difference between the two groups also increases with the number of children born in a single confinement. Twins are  $1\frac{1}{4}$  times as frequent for nonwhite as for white mothers, and quadruplets are  $4\frac{1}{2}$  times as frequent.

\* The frequency of plural births. Statist. Bull. Metrop. Life Insur. Co. 31:6-7, 1950.

born in Canada in 1934, and the Diligenti quintuplets, born in Argentina, 1943.

Quadruplets are born only once in 620,000 confinements. One set of triplets appears in every 9,400 births; twins are born once in every 92 confinements.

## Parenteral Fluid Therapy for Children

ALBERT M. HAND, M.D., AND C. R. LEININGER, M.D.\*

*Children's Memorial Hospital, Chicago*

**S**ERIOUS derangement of water and electrolyte balance frequently accompanies children's diseases.

Acute diarrhea, diabetic acidosis, pyloric stenosis, severe burns, and other conditions require parenteral replacement of water and some electrolytes. Successful therapy entails a knowledge of the type and approximate amount of electrolytes lost, explain Albert M. Hand, M.D., and C. R. Leininger, M.D.

In addition, the regular daily requirements of the child must be met (Table 1). Finally, the composition of the solutions employed in fluid therapy should be known (Table 2).

Serum electrolyte concentrations are best expressed in milliequivalents, mEq. per liter. Other terms, such as milligrams or volumes per cent, are chemically unrelated and therefore misleading. Milliequivalents are calculated by dividing the weight of the dissolved electrolyte, in milligrams per liter, by the atomic weight and multiplying by the valence.

For example, physiologic saline solution contains 0.9 gm. of sodium chloride per 100 cc. In 1 liter is 9 gm., or 9,000 mg., of sodium chloride. The atomic weight of sodium is 23; of chloride, 35.4. The va-

TABLE 1. BASIC DAILY REQUIREMENTS

Age	Weight kg.	Water cc./kg.	NaCl gm.	Protein gm./kg.	Calories/kg.
<1 yr.	2-10	150	0.5-1	1.5	60
1-2 yr.	10-13	120	1.5	1.0	55
4 yr.	15-20	80	2	1.0	40
10 yr.	30-40	50	3-4	0.6	30
>10 yr.	30+	45-30	6	0.6	25

Fluids are best given intravenously or subcutaneously with hyaluronidase to facilitate absorption. Hypertonic solutions and amigen should be administered only intravenously.

If blood is given, the rate is regulated to 10 to 20 drops per minute or 30 to 60 cc. per hour. In a single infusion, 20 to 30 cc. of blood per kilogram of body weight is the most that should be given to a child.

lence of sodium and chloride is 1. Therefore  $(9,000 \div 58.4) \times 1 = 154$  mEq. of sodium and chloride per liter.

The electrolyte pattern of the serum may be roughly calculated by knowing only the serum sodium-chloride and serum carbon-dioxide content. Serum carbon-dioxide expressed in volumes per cent may readily be converted to milliequiva-

\* Parenteral fluid therapy in children. *M. Clin. North America* 34:53-70, 1950.

TABLE 2. EQUIVALENTS OF REPLACEMENT SOLUTIONS

100 cc. of ↓	→ are equivalent to		
	NaCl gm.	Protein gm.	Calories
Physiologic saline	0.9		
Plasma or serum	0.45	5	20
Whole blood	0.45	6	24
5% amigen with 5% dextrose in distilled water	0.20	4	36
5% dextrose in distilled water			20
5% dextrose in physiologic saline	0.9		20
10% dextrose in distilled water			40

lents per liter by dividing by the factor 2.2.

The normal serum carbon dioxide is 27 mEq. per liter, 60 volumes per cent  $\div$  2.2. An infant weighing 10 kg. has approximately 7 liters of fluid, since 70% of the body weight is fluid. If acidosis is present with a serum carbon dioxide of 14 mEq. per liter, an alkali deficit exists of (27-14)  $13 \times 7 = 91$  mEq.

To correct this deficit, 91 cc. of one molar sodium lactate is required since each cubic centimeter of a univalent molar solution contains 1 mEq. This must be given as the isotonic one-sixth molar sodium lactate solution, so (91  $\times$  6) 546 cc. is required.

Roughly, 4.2 cc. of one-sixth molar sodium lactate solution per kilogram of body weight will raise the serum carbon dioxide 1 mEq. Probably only one-half of a calculated deficit should be replaced at one time.

Acute diarrhea causes a loss of water and base, principally sodium. Potassium, calcium, and magnesium are also lost. Metabolic acidosis results. The urine is acid in reaction, the blood pH lowered, and Kussmaul breathing will begin if the serum carbon-dioxide content is 15 mEq. per liter or less.

Therapy is aimed at replacing base and fluid. Hypotonic saline and one-sixth molar sodium lactate solution may be used. During recovery, calcium may be required.

Diabetic acidosis is another example of metabolic acidosis. Therapy should start with an intravenous infusion of 2 parts normal saline and 1 part one-sixth molar sodium lactate solution. The fluid deficit of a child in diabetic coma is usually equal to 5 to 10% of the body weight. This deficit should be replaced within the first twenty-four hours in addition to the normal daily fluid requirement. Initially, 1 unit of crystalline insulin per kilogram of body weight is given intravenously and the same amount subcutaneously.

Pyloric stenosis with vomiting causes metabolic alkalosis. The urine is alkaline, the blood pH elevated, and respirations slow or apneic. Acidification is indicated.

One-sixth molar ammonium chloride solution can be employed. For mild alkalosis, 15 cc. of one-sixth molar ammonium chloride is given in physiologic saline per kilogram of body weight. An additional 15 cc. of physiologic saline per kilogram is administered.



## Perforation of the Esophagus

JOHN D. KERNAN, M.D.\*

*Presbyterian Hospital, New York City*

A GRAVE surgical emergency is presented by perforation of the muscular and fibrous layers of the esophagus.

Leakage into the mediastinum results in periesophageal cellulitis with rapid spread of infection through the surrounding loose connective tissue. Until the advent of chemotherapy, antibiotics, and modern chest surgery, the condition was almost invariably fatal.

Symptoms given by John D. Kernan, M.D., are:

1] Pain and difficulty in swallowing

2] Exquisite tenderness in the lower part of the neck associated with crepitus

3] Rapid increase in area and severity of these signs with continuous pain not relieved by sedation in chest, back, or abdomen and, sometimes, hematemesis and nausea

4] Effusion in the posterior mediastinum with possible collapse of one or both lungs, seen on roentgenograms as widening of the mediastinal area and compression of the lungs.

5] Board-like rigidity of the abdomen

Perforation may result from internal or external trauma, disease, or congenital defect or, rarely, may be spontaneous.

The most frequent cause is in-

ternal trauma by endoscopic procedures for removal of foreign bodies, biopsy, or dilatation of strictures.

Removal of impacted foreign bodies is never a wholly safe procedure. Prompt hospitalization and careful roentgenographic diagnosis are essential. Intratracheal general anesthesia is necessary for adequate relaxation. Endoscopic examination should be done only by an experienced operator, and with gentleness.

Rough, sharp objects such as splinters of chicken or fish bone and open safety pins should first be located by a biplane fluoroscope and then removed with the appropriate tube and forceps. Coins and similar objects may be retained without harm for some time, unless they block the lumen of the esophagus so as to lead to aspiration of food and resultant bronchopneumonia.

The ingestion of corrosives like lye may cause sloughing of the esophageal wall with secondary perforations. Once the immediate emergency is past, the lumen may be kept open by daily dilation for twelve to twenty-five weeks with a mercury-weighted bougie, which is left in place for thirty minutes each time, or by gastrostomy and retrograde bougienage over a string or filiform guide preferably with fluoroscopic observation.

\* Perforation of the esophagus as a surgical emergency. *S. Clin. North America* 30:405-419. 1950.



Stricture after external trauma such as stab wounds is treated by dilation or secondary resection.

Perforation from chronic diseases, mainly cancer or tuberculosis, usually does not cause acute illness. The slow penetration of the esophageal wall is associated with formation of a protective zone of reaction.

Spontaneous rupture is rare, and results from gluttony, alcoholism, persistent vomiting of pregnancy or seasickness, epilepsy, or sudden pressure changes. Death is prevented only by prompt opening of the chest. The stomach contents are flushed out and the perforation closed by suture. The chest is then drained.

## Diabetic Retinopathy

BRUCE L. KANTAR, M.D.\*

THE earliest and most typical diabetic lesions in the retina are punctate red spots occurring with normal arterioles. These minute lesions, long considered hemorrhagic, are now known to be capillary aneurysms.

On reviewing 49 cases of diabetic retinopathy at the University of Minnesota Hospitals, Minneapolis, Bruce L. Kantar, M.D., noted microaneurysms in 48, with no other changes in 7. Fundi in the exceptional instance were obscured by dense retinitis proliferans. Subjects were twenty-two to seventy-eight years old and had been diabetic one to twenty-six years.

The punctate spots appear in the posterior pole of the eye, usually near the macula, and involve the venous end of the retinal capillary nets. Diameter varies from 30 to 90 microns and may be 15 times the width of the parent capillary.

Some aneurysms undergo thrombosis and are seen as round white nodules, others rupture and cause deep hemorrhages. The central retinal area may also contain small yellowish white exudates, single or clustered.

Larger retinal veins may become overfull and tortuous, with segments of pronounced dilatation and constriction, exudate along the walls, and extensive bleeding. In some cases, a network of new vessels arises in a brushlike mass supported by delicate connective tissue.

If hypertension develops, arteriosclerosis is shown by cotton-wool patches and focal arteriolar narrowing. Kimmelstiel-Wilson disease should be suspected when longstanding diabetes is associated with hypertension, renal insufficiency, and advanced retinopathy.


\* Diabetic retinopathy. *Bull. Univ. Minnesota Hosp.* 21:352-360, 1950.

# Femoral Hernia

F. M. AL AKL, M.D.


Kings County Hospital, New York

KEEP THIS PICTURE IN MIND




KEEP THIS PICTURE IN MIND


## SURGICAL TECHNIGRAM




1. Incise skin obliquely outward beginning 1 cm. medial to pubic tubercle, then extend incision half-way to anterior superior iliac spine.




2. Part and lift skin with sharp retractors. Incise the fat and superficial fascia. Clamp branches of external pudic and superficial epigastric vessels.




3. Open wound lateral to hernial mass and continue incision down to underlying inguinal ligament. Lift the lateral angle of wound.



4. Sweep fat and areolar tissue over inguinal ligament and around hernial protrusion. Isolate herniated mass and clear of adherent cribriform fascia. Clamp and cut branches of long saphenous vein.




5. Lift hernial mass and dissect neck free, exposing falciform ligament laterally, emerging saphenous vein caudad, and pectineous fascia medially. Tie bleeders.




6. Clamp, then incise longitudinally outer covering of ballooned-out cribriform fascia and crural septum of transversalis fascia with intervening and underlying fat down to peritoneal sac.



7. Pick up peritoneum and cautiously incise between forceps. Clamp the edges.




8. Enlarge opening, examine, then gently drag contents into sac to free and expose constricted portion at neck.




9. Reduce contents. Pull out on sac; transfix and tie neck flush.

## SURGICAL TECHNIGRAM



10. Excise sac and push stump back beyond femoral ring into abdominal cavity.




11. Doubly ligate and cut saphenous vein, then approximate falciform ligament to pectineous fascia and close skin.




12. If hernial contents are not reducible, retract skin cephalad and expose external inguinal ring and adjoining external oblique aponeurosis.




13. Split external oblique aponeurosis over cord as for inguinal hernia; then clamp and retract flaps.




14. Expose medial half of shelving edge of inguinal ligament, retract cremaster layer with underlying round ligament cephalad, then scissor cremaster attachment to inguinal ligament.




15. Retract round ligament with overlying cremaster from inguinal ligament. Incise exposed fascia from pubic tubercle to deep epigastric vessels parallel to and  $\frac{1}{4}$  cm. above inguinal ligament.




16. Insert finger into adipoareolar fossa and isolate neck of sac as it passes into femoral canal.




17. Retract transversalis fascia cephalad exposing hernial sleeve. Look for an anomalous obturator artery at the medial border of the femoral ring. Ligate and cut if present.




18. Insert grooved director into medial aspect of femoral canal and incise lacunar ligament medially over grooved director.




19. Reduce the herniated structures through the enlarged femoral ring; release clamp and pull sac out of femoral canal changing femoral into direct inguinal hernia.




20. Apply clamps to edge of sac, then purse-string and ligate neck of sac. Excise sac and cut ligature.




21. Depress floor of inguinal fossa, then approximate medial cut edge of transversalis fascia to pubic ligament.




22. (Left) Continue outward, push femoral sheath laterally with finger as final one or two sutures between transversalis fascia and pubic ligament are placed.



23. (Left) Suture conjoined structures to inguinal ligament.



24. (Right) Reposit round ligament and reconstruct external ring by suturing cruri together; then approximate balance of external oblique flaps.



25. (Right) Approximate skin with vertical mattress sutures.

## NOTES

Because of a wider femoral ring, femoral hernia occurs 6 times oftener in females than in males, and owing to the more rigid boundaries of the femoral ring, the incidence of strangulation is much higher with femoral than with inguinal hernia.

Diagnosis of femoral hernia may be difficult. The femoral mass may slip up and then ride over the inguinal ligament and thus be mistaken for an inguinal hernia. Inguinal adenitis with pain, nausea, vomiting, and cough impulse has been mis-

## SURGICAL TECHNIGRAM

taken for strangulated femoral hernia and vice versa.

Many authorities describe separate femoral and inguinal incisions for the repair of femoral hernia. The unpredictability of the findings and the difficulty of reducing the contents permit no such empiricism. An incision through which the entire inguinofemoral region can be reached should be used routinely.

Section of the lacunar ligament does not always permit reduction of the herniated structures. In such cases the incision into the lacunar ligament may be carried up into the inguinal ligament in Z fashion, or over the pubic tubercle cutting thereby the pubic insertion of the inguinal ligament. After the femoral ring is opened widely and the sac and contents are properly handled, the cut ends of the inguinal ligament are sutured together, or the severed ligament is sutured down to the pubic ligament together with the medial edge of the transversalis fascia during the repair.

Femoral hernias are covered with much fat, each layer of which may look like omentum. The texture of omental fat is distinct, however, and the omentum is usually preceded by the escape of the transudate in the sac. Strangulated or irreducible omentum may be segmentally ligated and excised. When doing so, the proximity of the transverse colon should be kept in mind and every clamp or suture should be applied with the assumption that the transverse colon may be just beyond.

Ordinarily the obturator artery arises from the internal iliac artery.

In approximately 25% of specimens it arises from the external iliac artery instead, either directly or from a common trunk with the deep epigastric artery. It then swings above the femoral vein to descend to the obturator foramen. Thus a hernial mass may push it to the medial or to the lateral aspect of the femoral ring. The artery may therefore be injured in manipulations around the femoral ring or section of the lacunar ligament. Likewise, both the pubic branch coursing along the inguinal ligament and the external spermatic branch of the deep epigastric artery may cause bleeding. The external spermatic branch may pierce the transversalis fascia medial to the cord and thus become injured during the dissection.

The proximity of the femoral vein both within the femoral sheath lateral to the femoral ring, and in the thigh behind the superior and inferior cruri of the falciform margin of the fascia lata, must be kept clearly in mind during the repair. Should the edge of the vein be punctured inadvertently during the repair, the bleeding point should not be clamped. The suture should be tied, not pulled out, and the bleeding will stop.

The function of sutures, here as elsewhere, is to hold surfaces together until fibroblasts grow between. Tissues are strangulated and destroyed by gross handling with bulky suture material and big bites. The bite of the needle need be no deeper, and the weight of the suture material no heavier than required to approximate adjoining surfaces.





**Stops  
wracking  
cough...**



**but keeps  
the cough  
reflex**

MERCODOL provides prompt, selective relief that doesn't interfere with the cough reflex needed to keep throat passages and bronchioles clear.

This complete, pleasant-tasting prescription contains a *selective* cough-controlling narcotic<sup>1</sup> that doesn't impair the beneficial cough reflex . . . an effective bronchodilator<sup>2</sup> to relax plugged bronchioles . . . an expectorant<sup>3</sup> to liquefy secretions. Remarkably free from nausea, constipation, retention of sputum, and cardiovascular or nervous stimulation.

# MERCODOL®

**THE ANTITUSSIVE SYRUP THAT CONTROLS COUGH—KEEPS THE COUGH REFLEX**  
An exempt narcotic

**MERCODOL with DECAPRYN®**  
for the cough with a  
specific allergic basis.



Each 30 cc. contains:

<sup>1</sup> Mercodione®	10.0 mg.
<sup>2</sup> Nethamine® Hydrochloride	0.1 Gm.
<sup>3</sup> Sodium Citrate	1.2 Gm.

# Medical Forum

*Discussion of articles published in MODERN MEDICINE is always welcome. Address all communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.*

## Non-Union of the Medial Malleolus\*

TO THE EDITORS: We are in complete agreement with the comments of Dr. Sam W. Banks on the treatment of nonunion of fracture of the medial malleolus. In our opinion one seldom finds a fibrous union sufficiently strong to guard against some pain and swelling of the ankle on a normal routine.

Screw fixation is the only thing we have found that gives adequate stability, and cancellous bone is preferable to cortical bone in promoting union. It is possible that the percentage of nonunions would be materially decreased by adequate primary reduction of the fragment even if primary reduction meant open operation and screw fixation at the time of original treatment.

R. G. TOWNSEND, M.D.  
Calgary, Alb.

► TO THE EDITORS: The method described by Dr. Sam W. Banks seems to be excellent and quite simple to perform. It is really a modification of various methods that have been published from time to time. Its simplicity recommends it.

\*MODERN MEDICINE, Oct. 1, 1949, p. 73.

I believe, however, that when the fracture of the malleolus involves the joint, the symptoms of which the patient complains and the signs noted on physical examination, such as Dr. Banks mentions, are usually from an associated traumatic arthritis resulting from damage to the ankle joint. The healing of the fracture will, in my opinion, not alleviate these symptoms, and I think the operation should be kept entirely for cases with definite mobility of the fragment. Certainly a bone graft of the internal malleolus, such as described, is not going to relieve the accompanying arthritis of the ankle joint.

H. J. SPOONER, M.D.  
Regina, Sask.

► TO THE EDITORS: The article by Dr. Sam W. Banks is precise and well illustrated.

The method described, which has been used for some years for painful nonunion of the medial malleolus, gives excellent results.

I believe that there is no better technic to advise for treatment of these cases.

LIONEL GROLEAU, M.D.  
Sherbrooke, Que.



An illustration of a Protamide product box and a syringe. The box is dark with white text and features two oval labels. The top label reads 'List No. 500', 'PROTAMIDE®', and '1.3 cc. Size'. The bottom label reads 'List No. 500', 'PROTAMIDE', and '1.3 cc. Size'. The box also has text for 'SHERMAN LABORATORIES', 'G. H. Sherman, M.D., Founder', 'DETROIT 15, MICHIGAN', and 'Wander, Cal., Los Angeles, Calif.'. A syringe is shown to the left of the box, with a needle pointing towards the top left.

# PROTAMIDE

SHERMAN

List No. 500  
**PROTAMIDE®**  
1.3 cc. Size

SHERMAN LABORATORIES  
G. H. Sherman, M.D., Founder  
DETROIT 15, MICHIGAN  
Wander, Cal., Los Angeles, Calif.

List No. 500  
**PROTAMIDE**  
1.3 cc. Size

## NEW, DRAMATIC THERAPY FOR THE RELIEF OF PAIN AND LESIONS OF **HERPES ZOSTER**

**DESCRIPTION:** Protamide is a sterile, aqueous colloidal solution of a specially processed proteolytic enzyme, for the maximum relief of nerve root pains of Herpes Zoster and Tabes Dorsalis.

**CLINICAL RESULTS:** Highly gratifying clinical results have been obtained with the use of Protamide (Sherman) in the treatment of the extremely resistant herpes syndrome. Pain has been relieved in the great majority of herpes cases within four to forty-eight hours and lesions have healed in ten days or less—regardless of the particular nerve roots involved. Complete clinical data may be obtained by writing for the Protamide literature on Herpes Zoster and a recent reprint on Protamide for Tabes Dorsalis.

**DOSAGE:** In Herpes Zoster the recommended dosage is 1.3 cc of Protamide intramuscularly each day from two to four days. Causes no reactions—comparatively painless—no contraindications or incompatibility. All Protamide is clinically tested for positive results. Can be stored at room temperature without loss of potency.

**SHERMAN LABORATORIES**  
G. H. Sherman, M. D., Founder  
**BIOLOGICALS • PHARMACEUTICALS**  
DETROIT 15, MICHIGAN

**REGISTERED U. S. TRADE MARK**

# Diagnostix

*Here are diagnostic challenges presented as they confront the consultant from the first clue to the pathologic report. Diagnosis from the Clue requires unusual acumen and luck; from Part II, perspicacity; from Part III, discernment.*

## Case MM-172

### THE CLUE

ATTENDING M.D.: The patient I'd like you to see today was in good health until a month ago, when he began to have diarrhea, six to ten loose stool movements a day without blood or cramps, and malaise. He worked for a week and then consulted a doctor, who treated him symptomatically. The doctor, on the note requesting admission to the hospital, said that the man's temperature was between 101 and 103° each day and that a trial of penicillin had not brought the fever down and was therefore discontinued. The patient is sixty-

eight years old. He has just been admitted to the hospital.

VISITING M.D.: (*Walking into the room and examining the patient*) He appears quite sick. I note his temperature is 103°, pulse 90; he has a maculopapular rash, some of the lesions are as large as 2 cm. on the abdomen and back. The liver is palpable but I cannot feel the spleen. Better get the routine laboratory work, including blood cultures. What do you think the man has?

ATTENDING M.D.: Well, I noted a systolic apical murmur, which you didn't mention. I'm inclined to think that he has subacute bacterial endocarditis.


VISITING M.D.: But the disease began with diarrhea.

ATTENDING M.D.: I know. Of course, we have no good evidence as yet of subacute bacterial endocarditis, but I intended to give him large doses of penicillin at once.

VISITING M.D.: Better get several blood, urine, and stool cultures first.

(Continued on page 82)





Psoriasis of 2½ years' duration.



Same case after 2 months' treatment with Mazon.


*Remission of the psoriatic lesion  
evaluates the effectiveness of* **MAZON**

- It is the marked capriciousness of psoriasis which makes difficult the evaluation of any method of treatment, but dermatologists agree that the most satisfactory approach to an effective regimen usually commences with local therapy.
- When systemic or metabolic involvement is not manifested—even though the condition is generalized or stubbornly resistant—Mazon has demonstrated its value in arresting psoriatic lesions and promoting symptomatic relief.
- Mazon is a compound of mercury salicylate  $\frac{1}{3}$  gr. to the ounce; benzoic acid; sodium stearate; salicylic acid and tars.


BELMONT LABORATORIES, Philadelphia, Pa.

**MAZON**


# FOR all TYPES



TRICHOMONAS VAGINALIS  
pH 5 to 6



MIXED INFECTIONS  
(Staphylococcus, Streptococcus,  
Escherichia coli, etc.)  
pH 5.8 to 7.8



MONILIA ALBICANS  
pH 5.5 to 6.8

Floraquin restores the normal flora.

SEARLE

RESEARCH IN THE SERVICE OF MEDICINE

# OF VAGINITIS...


By reestablishing a normal epithelial glycogen content, acidity and Döderlein bacilli, Floraquin may be described as the complete restorative treatment in vaginitis.

**FLORAQUIN®**—a product of Searle Research—combines the potent trichomonacide, Diodoquin-Searle, with lactose, dextrose and boric acid.

Floraquin Powder—for office insufflation.


Floraquin Tablets—for patient's use.

**G. D. SEARLE & CO., Chicago 80, Illinois**



**DÖDERLEIN BACILLI**  
pH 3.8 to 4.4

and normal  
vaginal mucosa



## DIAGNOSTIX

### PART II

VISITING M.D.: Do you know of epidemic diarrhea in town?

ATTENDING M.D.: I haven't heard of any.

VISITING M.D.: Let's call the Department of Health and ask them. *(The call is made. The Department of Health has no reports at present of any similar infectious gastrointestinal diseases.)*

VISITING M.D.: The patient's abdomen is quite distended. I think you should give him a milk diet. He has not vomited. I will talk to him some more. *(Long interview with the patient)* The patient tells me that he had fever for two or three days at the onset, then he felt well for about four or five days. The fever had a brisk recurrence and has been up and down ever since. Whatever the man has, it is a relapsing illness,

or at least it has had a significant relapse. And yet the signs aren't those of a relapsing fever.

ATTENDING M.D.: No.

VISITING M.D.: You'd better put a Wangenstein tube down and start suction.

### PART III

ATTENDING M.D.: *(Two days later)* The blood and stool cultures both came back positive for *Eberthella typhosa*; we have started Chloromycetin. The temperature today is 105°, pulse 120. He has numerous petechial hemorrhages. White count is normal. The abdominal distention has diminished considerably; we have removed the tube.

### PART IV

VISITING M.D.: Just because we don't see much typhoid fever is no reason why we should not be very much on the alert for it. I'm glad to have this opportunity of discussing typhoid fever, which has practically disappeared from the state. We must get the epidemiologist to investigate this case and discover a possible carrier. Most likely the patient contracted the disease from some carrier. His attack has been a relatively mild one. Typhoid fever used to run about three weeks before the patient began to improve. Sometimes it ran four to five weeks. The skin rash is obviously composed of the characteristic rose spots. I'm not surprised you didn't recognize them, since you had never seen them before. I saw them when I was an intern. We must watch out that the artificial controls over water




"See here, Doctor, I'm in a hurry."

when  
temptation  
crashes  
the  
party

● Uninvited he comes, arousing a mutinous appetite, then destroying her every intention of following her diet. When these dietary misdemeanors become frequent, there couldn't be a better time to employ DESOXYN Hydrochloride. By its combined effect of depressing the appetite and elevating the mood, DESOXYN helps you encourage dietary adherence.

Weight for weight DESOXYN is more potent than other sympathomimetic amines. Thus small amounts produce the desired cerebral effect without producing undesired side-effects. One 2.5-mg. tablet before breakfast and another about an hour before lunch is usually sufficient. A third tablet may be taken in midafternoon if necessary and if it does not cause insomnia. DESOXYN's other advantages are faster action, longer effect. In small oral doses, no pressor effect has been observed. With proper dosage,

DESOXYN is safe, simple, effective. **Abbott**



note the name

**DESOXYN<sup>®</sup> hydrochloride**

(Methamphetamine Hydrochloride, Abbott)

**TABLETS**  
2.5 and 5 mg.

**ELIXIR**  
20 mg. per fluidounce  
(2.5 mg. per fluidrachm)

**AMPOULES**  
20 mg. per cc.





# RUTAMINAL\*

Ocular  
Fundus in  
Degenerative  
Vascular  
Disease—  
Hypertension,  
Diabetes,  
Arteriosclerosis  
—note  
tortuous  
blood vessels,  
areas of  
exudation,  
hemorrhagic  
areas



Normal  
Ocular  
Fundus



In keeping with newer clinical findings, the rutin content of RUTAMINAL has been increased to 60 mg. per tablet (three times the former rutin content) at no increase in cost to the patient.

\*RUTAMINAL is the trademark of Schenley Laboratories, Inc. and designates exclusively its brand of tablets containing rutin, aminophylline, and phenobarbital.

the  
protection  
of  
**rutin**<sup>1</sup>  
the  
action  
of  
**aminophylline**  
the  
sedation  
of  
**phenobarbital**  
—for  
use  
in  
selected  
cardiovascular  
and  
diabetic  
conditions  
in  
which  
excessive  
capillary  
fragility  
presents  
a  
complicating  
hazard  
—bottles  
of  
100  
tablets

schenley laboratories, inc., 350 fifth ave., new york 1, n. y.

© Schenley Laboratories, Inc.

and milk supply do not break down so that we have epidemics again. Although aureomycin and Chloromycetin have both been recommended for the treatment of typhoid fever, Chloromycetin definitely seems to be the drug of choice. The 3-gm. initial dose is followed by 2 gm. daily for about a week after symptoms have subsided. We'll still be concerned with the problem of carriers if Chloromycetin doesn't eradicate biliary typhoid foci. We will, of course, need three successive negative stools and urine cultures at intervals of at least one week, beginning one week after subsidence of clinical symptoms and after the drug has been stopped for a week. We should make sure that all the doctors and nurses in the hospital who have been exposed to the patient in any way

have adequate typhoid vaccination. Typhoid vaccine administered even after exposure to typhoid reduces the infection rate. Certainly the family should be vaccinated.

ATTENDING M.D.: Would you give Chloromycetin in divided doses?

VISITING M.D.: Yes.

ATTENDING M.D.: What dose would you give of the vaccine?

VISITING M.D.: For anyone who had had a full course of vaccine within the year, give 0.1 cc. intradermally. If the interval is longer, 0.5 cc. subcutaneously.

ATTENDING M.D.: Do you believe the patient may have a relapse after we stop the Chloromycetin?

VISITING M.D.: It is quite possible. We should continue the drug longer than seems necessary, since the exact effect of Chloromycetin with this relatively rare disease is not known.



"So you just got back from a vacation? Hmm. In that case I'd say you need a good long rest."

## Life's Weary Moments

Think of a gag that fits the illustration. For every issue a new gag is published and the author is sent \$5. The August 1 winner is

John Pellettieri, M.D.  
Cicero, Ill.

Mail your caption to  
The Cartoon Editor  
Caption Contest

No. 1

MODERN MEDICINE  
84 South 10th St.  
Minneapolis 3, Minn.


# WHAT THE BREAKFAST CEREAL CONTRIBUTES


## *in the Cereal and Milk Serving*

That breakfast cereals make a significant contribution to the over-all nutrient content of the cereal and milk serving is readily apparent from the bar chart. In the case of five of the nine nutrients, as well as in calories, the breakfast cereal alone provides more than 50 per cent of the total amounts contributed by both cereal and milk, and almost 50 per cent of the protein.

These figures are based on composite averages of all breakfast cereals—whole grain, enriched, or restored to whole grain values of thiamine, niacin and iron.

The values shown demonstrate that breakfast cereals are significant sources of many essential nutrients. Their inclusion in the balanced breakfast goes far in satisfying the daily requirements for the nutrients they contribute. For this reason they are widely recommended by nutritionists as an important component of breakfast.





The chart illustrates how the chief components of the cereal serving (1 oz. of breakfast cereal, 4 oz. of milk, 1 teaspoonful of sugar) nutritionally supplement each other.

In addition to contributing a host of essential nutrients, breakfast cereals perform another, largely unrealized, function. In 1948, an estimated two billion quarts of milk were consumed with breakfast cereals. This quantity represents 16 billion cereal servings, or 66 quarts of milk per family per year. In view of the fact that about 40 per cent of our adult population does not drink milk as such, breakfast cereals perform the outstanding function of bringing milk into the dietaries of many adults who otherwise would not receive this essential food. Thus they contribute to the calcium intake of many individuals and aid in overcoming the widespread calcium deficiency of many adult diets.\*

\*Stearns, G.: Human Requirement of Calcium, Phosphorus and Magnesium, Council Report, J.A.M.A. 142:478 (Feb. 18) 1950.



The presence of this seal indicates that all nutritional statements in this advertisement have been found acceptable by the Council on Foods and Nutrition of the American Medical Association.

## CEREAL INSTITUTE, INC.

135 South La Salle Street • Chicago 3

A RESEARCH AND EDUCATIONAL ENDEAVOR DEVOTED TO THE BETTERMENT OF NATIONAL NUTRITION

# Short Reports

## OBSTETRICS

### Complications of Late Pregnancy

Thromboplastin may be a chief activator of some diseases of late pregnancy. Dr. Charles L. Schneider of Wayne University, Detroit, finds that reactions similar to some complications of human pregnancy can be induced in rabbits by placental injury. These complications are apparently produced by intravascular coagulation caused by the presence in the maternal circulation of material from the damaged placenta. Evidence suggests that thromboplastin is probably the material which initiates the clotting. The most common result of the trauma, liver necrosis, closely parallels the reaction to injection of thromboplastin extracts.

*Surg., Gynec. & Obst.* 90:613-622, 1950.

## ANTIBIOTICS

### Cyclotron-purified Drugs

Antibiotics may be sterilized by a high-voltage atom-smashing machine, similar to the cyclotron employed for theoretic investigations. At the Upjohn Pharmaceutical Company, Kalamazoo, Mich., a 2,000,000-volt accelerator beams a 175,000-mile-per-second stream of electrons into tubes and bottles of penicillin and streptomycin to free the antibiotics of bacteria. Previously, heat, filtration, and chemical treatment had been necessary for sterilization.

## METABOLISM

### Thiourea Hyperglycemia

Oral administration of potassium iodate appears to protect rats against hyperglycemia induced by thiourea injections. In untreated animals injected with the goitrogenic drug, Dr. David E. Mann, Jr., of Purdue University, Lafayette, Ind., observed an average rise in blood sugar of 91 mg. per 100 cc. The same injections produced an increase of less than 15 mg. per 100 cc. in rats that received potassium iodate in drinking water.

*Proc. Soc. Exper. Biol. & Med.* 73:657-658, 1950.

## EXPERIMENTAL SURGERY

### Mitral Valve Prosthesis

Study of the effects of mitral valvular disease on cardiovascular and respiratory physiology is facilitated by use of an artificial mitral valve. A prosthetic valve designed by Dr. G. Rehmi Denton and co-workers of Albany Medical College, N.Y., can be substituted surgically for the mitral valve of dogs. The prosthesis, which has no moving parts, may be constructed so as to give varying degrees of mitral stenosis or regurgitation. Several dogs have lived for as long as five months with the artificial valve in place. The animals show no ill effects if the prosthesis is constructed to function as a normal valve.

*Federation Proc.* 9:31, 1950.



## 2 to 3 cc. **KOAGAMIN®**

**PREOPERATIVELY**—2 to 3 cc. of KOAGAMIN—prevents oozing, allows the surgeon a clearer field of operation and reduces the need for local hemostatic measures.  
**POSTOPERATIVELY**—2 to 3 cc. of KOAGAMIN—aids control of secondary bleeding.

**THERAPEUTICALLY**—2 to 3 cc. of KOAGAMIN—aids in control of bleeding in gastric and duodenal ulcers, hematemesis, hematuria, hemorrhagic purpura, epistaxis, blood dyscrasias, etc.

KOAGAMIN's prompt action—a matter of minutes—differs from that of vitamin K, which must first be converted to prothrombin in the liver—a matter of hours. Vitamin K is useful *only* in cases where prolonged prothrombin time is a factor. *Even in these cases,* KOAGAMIN should also be used for its rapid action.

*Supplied in 10 cc. diaphragm-stoppered vials.  
Write for comprehensive dosage chart and literature.*



**CHATHAM PHARMACEUTICALS, INC.**  
**NEWARK 2, NEW JERSEY, U. S. A.**

*Available Through Your  
Physician's Supply House or Pharmacist*

## Doctor to Doctor

Think of gag that fits the illustration. For every issue a new gag is published and the author is sent \$5. The August 1 winner is

Laird McNeel, M.D.  
Genoa City, Wis.

Mail your caption to  
The Cartoon Editor  
Caption Contest  
No. 2

MODERN MEDICINE  
84 South 10th St.  
Minneapolis 3, Minn.



"Well, John, are you sure you always wear the lead apron for fluoroscopy?"



# SWIFT...

**ESTIVIN** promptly relieves ocular and nasal symptoms caused by pollinosis or hay fever.

**ESTIVIN** is effective even in cases where antigen treatment and antihistamines have failed to relieve ocular and nasal irritation.

**ESTIVIN** used daily commonly relieves and soothes conjunctivitis and alleviates ocular distress such as itching, burning eyes and excessive lacrimation.

**ESTIVIN** does not cause secondary reactions.

Samples and literature on request.



## ESTIVIN

**Schieffelin & Co.**  
PHARMACEUTICAL AND RESEARCH LABORATORIES  
24 COOPER SQUARE • NEW YORK 3, N. Y.





The effective  
triple  
sulfonamide  
that eases  
your mind  
about possible  
toxic effects

Pansulfa offers you the most reliable safeguards against crystalluria and renal damage. This effective triple sulfonamide contains sulfacetamide—the least toxic sulfonamide studied.\* Your prescription for Pansulfa offers

- 1 The established antibacterial power of three sulfas.
- 2 Less danger of crystalluria or renal damage.
- 3 Uniform dosage—the thixotropic gel of the suspension assures even dispersion. Also available in palatable tablets.

Pleasant tasting

# PANSULFA

SULFACETAMIDE  
SULFADIAZINE  
SULFAMERAZINE

Each teaspoonful or tablet contains 0.5 Gm. (7½ grs.) of the rapidly soluble sulfonamides 1:1:1



\*See Lehr, D: Federation Proc. 8:315 (1949)

"PANSULFA" trade-mark

# Detailed Dosage Data for Physicians

A 20-page brochure prepared exclusively for the medical profession presents busy physicians with detailed dosage information relating to quantity, frequency and duration of administration in relation to menses, as well as indications, rationale, etc., regarding ERGOAPIOL (Smith) with SAVIN. This time-tested uterine tonic is thoroughly described in this brochure, "Menstrual Disorders—Their Significance and Symptomatic Treatment". A copy, available to physicians only, will be supplied on request. Ethical since its inception, ERGOAPIOL (Smith) with SAVIN is dispensed only on your prescription.

**INDICATIONS:** Amenorrhea, Dysmenorrhea, Menorrhagia, Metrorrhagia, and to aid involution of the postpartum uterus.

**GENERAL DOSAGE:** 1 to 2 capsules, 3 to 4 times daily—as indications warrant.


In ethical packages of 20 capsules each, bearing no directions.




Ethical protective mark, M.H.S., visible only when capsule is cut in half at seam.

**ERGOAPIOL (SMITH) WITH SAVIN**

Martin H. Smith Co. • 150 Lafayette St. • New York 13, N.Y.





*Critical diagnosis quicker  
with the Bausch & Lomb*  
**MAY OPHTHALMOSCOPE**

Illumination of the fundus is brilliant,  
colors are more natural, and a field free from  
filament images and spots is made possible with  
the new Daylight Blue Light and May prism.  
Lenses, fingertip controlled, range from +20  
to -25.00D with numbers magnified and  
illuminated for easy reading in dark.  
Adjustable rheostat in handle.  
Available in handsome, durable case.



# Washington Letter

## Geriatrics Is of First Concern to National Conference

The first National Conference on the Aging, which meets in Washington this month, will consider how to care for and how to make life satisfying for the aged and aging. The problems are new, as medical and social developments are reckoned, because until fifty years ago people rarely lived to old age and the shortcomings of a society that had little security or dignified comfort for the aged were largely unappreciated.

The conference will run for three

days, from August 13 to 15. According to Federal Security Administrator Oscar Ewing, whose agency is sponsoring the meeting, government officials have no blueprint, even tentative, for a solution. But they hope, Mr. Ewing says, that out of the conference will come a better understanding of the problem and better direction of effort.

To the conference have been invited lay and professional people

(Continued on page 98)



"Now we can cancel all those magazine subscriptions."

**Molybdenized Iron . . .**

**The Most Effective**

**Iron Therapy**

**Known . . .**

*White's* **Mol-iron®**

**NOW ALSO  
with LIVER AND VITAMINS**  
(including B<sub>12</sub>)

For iron-deficiency anemias associated with excessive drain on nutritional reserves—e.g., in chronic infection, malignancy, prolonged anorexia, post-surgical and other convalescent states.

**Mol-Iron with Liver and  
Vitamins provides, in each capsule:**

Mol-Iron . . . . .	198 mg.
Desiccated Whole Liver . . . . .	0.45 Gm.
(equivalent to 1.8 Gm. whole liver—minus water content only—not a fraction or an extract)	
Thiamine hydrochloride . . . . .	1 mg.
Riboflavin . . . . .	1 mg.
Vitamin B <sub>12</sub> . . . . .	1 µg.
Nicotinamide . . . . .	5 mg.
Pyridoxine hydrochloride . . . . .	0.5 mg.
Calcium pantothenate . . . . .	1 mg.

Supplied in bottles of 100 capsules. Average dose  
1 to 2 capsules three times daily after meals.

WHITE LABORATORIES, Inc., Pharmaceutical Manufacturers, Newark 7, N. J.

# PABALATE®

*with demonstrated*



Technique of


Application of

Medication

Medication



**CLINICAL  
EFFICACY**



LIDUM VALUATE 72% PARALATE 2.4

Percentage of patients in whom relief was obtained in 24 hours

**PROLONGED  
(24 HOUR)  
RELIEF**



**COMPLETE  
FREEDOM  
FROM  
TOXICITY**

LIDUM VALUATE 55.2% PARALATE 0

Percentage of patients in whom relief was obtained in 24 hours

Percentage of patients in whom relief was obtained in 24 hours



# antirheumatic superiority...

Experience in the administration of the antirheumatic Pabalate confirms the efficacy, reliability and safety\* of its synergistic combination of salicylate and para-aminobenzoic acid.<sup>1,2,3,4,5</sup> Pabalate has been reported not only to provide "twenty-four hour pain relief,"<sup>6</sup> but its use (unlike that of salicylate alone) carries a high degree of freedom from toxic reactions.<sup>6</sup>

**INDICATIONS** rheumatoid arthritis, fibrositis, acute rheumatic fever, gout, osteoarthritis. The Liquid is also recommended as a replacement for analgesic-antipyretic medication generally.

**FORMULA** each Pabalate Tablet or each 5 cc. (one teaspoonful) of Pabalate Liquid contains sodium salicylate, U.S.P. (5 grs.) 0.3 Gm., para-aminobenzoic acid (as the sodium salt) (5 grs.) 0.3 Gm.

**SUPPLIED** Pabalate Tablets are supplied in bottles of 100 and 500; Pabalate Liquid in pints and gallons.

## REFERENCES

1. Beckman, H.: *Treatment in General Practice*, 6th ed., W. B. Saunders Co., Phila., 1948.
2. Belisle, M.: *Union méd. du Canada*, 77:392, 1948.
3. Dry, T. J. et al.: *Proc. Staff Meet., Mayo Clin.*, 21:497, 1946.
4. Rosenblum, H. and Fraser, L. E.: *Proc. Soc. Exper. Biol. & Med.*, 65:178, 1947.
5. Salassa, R. M., et al., *J. Lab. & Clin. Med.*, 33:1393, 1948.
6. Smith, R. T.: *J. Lancet*, 70:192, 1950.


**A. H. ROBINS CO., INC., Richmond 20, Va.**

*Ethical Pharmaceuticals of Merit since 1878*

# PABALATE



\*The data for these graphs are derived from the tabular material accompanying "Treatment of Rheumatoid Arthritis and other Rheumatic Conditions with Salicylate and Para-Aminobenzoic Acid: a study of 125 patients." by Richard T. Smith, *J. Lancet*, 70:192, 1950.





## WASHINGTON LETTER

from every phase of American life concerned with the aged—from educational, industrial, fraternal, civic, labor, and medical organizations. However, invitations went out to individuals, not to organizations.

More than 1,200 invitations were sent, and acceptances to date indicate that about 1,000 persons will attend. Work sessions or discussion groups are scheduled in each of the following fields: population changes, employment and rehabilitation, financial income, health maintenance and service, education, family life and housing, recreation, religion, community organization, and the training of personnel for working with the aged.

FSA officials are planning for a large representation from various fields of medicine, although no organization will be represented as such. Mr. Ewing said that his only thought on the health question is that the conference will result in better appreciation of what specific medical knowledge should be applied to old age. He also hopes to eliminate some duplication of medical research in geriatrics and to facilitate exchange of information.

In a background report on the aging population, FSA lists the major objectives as control of chronic disease, research into the aging process, development and instruction in rules for living with the infirmities and illnesses of old age, and plans for the care of ill and infirm old people. Discussions at the conference will center around these points.

In this report FSA states:

Crippling and disabling illnesses act as a drain both on the individual and

on society by preventing gainful employment, by setting up a group of people who are dependent on others for their livelihood and care by necessitating vast networks of nursing homes and specialized institutions.


In the same report, FSA strongly recommended adoption of a system of federal insurance for the temporary and permanently disabled.

Attention of the medical profession is directed toward research on how, why, and what makes people age and the manner of deterioration of function of tissues, muscles, and organs.

Most physicians are aware, in a general sense, that treatment of the aged was not a pressing problem before 1900 and that old people did not then constitute the social and economic burden they do now. FSA has presented some statistical information on the change. In the fifty years before 1900, life expectancy had increased only seven years, from forty to forty-seven. In 1900 almost 1 in every 2 persons was under twenty years of age, and only 1 in 25 had reached sixty-five years. By 1940, the proportion of children and youth had decreased to 1 in every 3 persons, and the proportion of the aged (sixty-five) had increased from 1 in 25 to 1 in 14. This year the ratio will be about 1 in every 13 persons. Expressed in percentages, in 1900 the aged constituted 4.1% of our population; this year they will constitute about 7.6%.

### Taft Stands Pat

An off-the-record visit by Oscar Ewing to Sen. Robert Taft (Rep., Ohio) was part of the administration's strategy on Reorganization



# now Feojectin

a completely new form of iron therapy



"We can now treat successfully those refractory iron-deficiency anemias that have previously defied us."

(Editorial, The Lancet, Jan. 1, 1949)

Feojectin is a stable solution of saccharated iron oxide for *intravenous injection only*. It is particularly indicated for those cases of iron-deficiency anemia in which oral medication (1) is ineffective, (2) is not well tolerated, or (3) produces results too slowly.

Feojectin is supplied in boxes of six 5 cc. ampuls. (Each ampul contains the equivalent of 100 mg. of elemental iron.)

## Feojectin

for use when oral iron fails

*Smith, Kline & French Laboratories  
Philadelphia*

"Feojectin" Trademark

## WASHINGTON LETTER

Plan No. 27, which would make FSA into a department of Health, Education, and Security. A similar proposal was made last year, only to lose out when Senate voted it down. Last year, the idea, because it came as a surprise to most House and Senate members, aroused needless opposition.

This time President Truman himself saw some of the senators, and Budget Bureau officials talked the idea over with others from both parties; Mr. Ewing volunteered to try to convince Sen. Taft. Neither Sen. Taft nor Mr. Ewing disclosed what was said at their meeting. However, Sen. Taft was not swung over.

A week or so after talking with Mr. Ewing, the Senator announced that he would oppose Plan 27 for the same reasons that he opposed the proposal last year. He objected because health activities would be further entangled with education and social security matters, rather than set up in an independent department or agency. Sen. Taft also said that he was fearful that all the top level thinking in the proposed new agency would be influenced by social rather than medical factors.

Late in the session Rep. Charles A. Wolverton (Rep., N. J.) introduced a bill for "reinsuring" health insurance plans, a provocative suggestion which served to get down in writing some of the ideas that had been advanced by, among others, Harold Stassen. Briefly, the bill proposed to set up a federal corporation that would protect insurance plans against catastrophic illnesses.

Here are the major provisions: Insured groups would pay 2% of

their gross premiums to the corporation. The corporation would have to meet the cost of an illness up to \$1,000 in any one year. After \$1,000, the corporation would pay two-thirds of all costs but the individual insurance plan would have to pay the remaining third. For hospital costs, the patient himself would be required to pay \$1 a day. The bill is carefully worded to rule out private-profit insurance plans.

The bill does not offer protection against a wave of high-cost cases, all under \$1,000, which have proved disastrous to many small nonprofit companies. Instead, a system is designed to limit the corporation's liability in case a company goes broke. If this happens, the corporation remains liable to an individual policy holder for the same benefits it would have paid the defunct association, but the corporation's liability ceases after one year. Mr. Wolverton's bill is attracting considerable interest from groups which want to see a wide extension of medical benefits, yet are opposed to national health insurance as represented in the administration's Thomas-Murray-Dingell bill.

### GI Medical Education

A ruling by Veterans Administration has ended an argument over extending the time in which veterans will be eligible for medical education. The deadline now is July 25, 1951. If the GI hasn't started his course by that time, he is no longer entitled to educational benefits.

On the assumption that a large number of qualified veterans would be unable to enter crowded medical

**MAXIMUM  
THERAPEUTIC  
RESULTS**

*prescribe*  
**NEOCYLATE**  
*the original*

TRADEMARK

**Salicylate-PABA Combination**

AVAILABLE IN TWO STRENGTHS  
REGULAR FORMULATION

THIS FORM AND STRENGTH

*Indications:* Effective against  
fever, pain, inflammation, and  
all types of rheumatism. Also  
effective in the treatment of  
menstrual pain.

*Precautions:* Do not use in  
patients with known hypersensitivity  
to salicylates or PABA. Do not  
use in patients with severe  
renal or hepatic disease.

**NONALCOHOLIC, HYPOALLERGENIC,  
FRUIT-FLAVORED SYRUP.**

*Warnings:* Do not use in patients  
with severe renal or hepatic disease.  
Do not use in patients with  
known hypersensitivity to salicylates or PABA.

**THE CENTRAL PHARMACEUTICAL CO.**  
P.O. Box 100, New York, N.Y. 10001  
NEW YORK - NEW YORK

## WASHINGTON LETTER

schools by that date, H.R. 7235 proposed extending the deadline. At a hearing on the bill, it was brought out that Budget Bureau officials were firmly opposed to the new deadline. They argued that this deadline was carefully selected as one which would give the GI every reasonable opportunity to start his medical course.

VA finally settled the argument by making the following ruling: Any eligible veteran who starts his premed course by next July 25 but is delayed in entering medical school by conditions beyond his control will be considered eligible for medical school benefits at any future date he may gain admission to medical schools. VA will be the judge of whether the student has made en-

ergetic efforts to gain admission to a school.

VA anticipates that only a relatively few veterans will fall in this class.

### Washington Notes

A special committee appointed to advise President Truman on VA hospital problems and requirements may have another and more important role. Mr. Truman will be able to refer to it any VA construction legislation which might be unnecessary or purely political in conception. This way, if the committee recommends a presidential veto Mr. Truman will not have to stand up alone against criticism, as he was obliged to do early last year when he cut



## START YOUR PRACTICE WITH THE RIGHT OFFICE RECORDS

### the **DAILY LOG**

#### Solves All Your Bookkeeping Problems

A complete financial record book for physicians—*designed by a physician*. Reduces billing mixups—helps increase your income by catching all charges due—*aids in keeping costs in line* by giving you an itemized account of all your expenses—*gives you all records needed for income tax return.*

#### Introductory Offer!

As a special "get acquainted" offer to doctors just beginning practice, the Dr. Colwell DAILY LOG for 1950 is offered at a reduced rate. You can use this proved system for the remainder of the year at considerable saving. WRITE for details and descriptive booklet.



- COMPLETE IN ONE VOLUME
- COSTS LESS THAN 2¢ PER DAY

**COLWELL PUBLISHING COMPANY**  
239 University Ave.      **Champaign, Illinois**



## RELAX THAT SPASM

with  
MESOPIN

When pain, heartburn,  
belching, nausea, or  
unstable colon are due to  
gastrointestinal spasm,

Mesopin provides an effective means for prompt relief. Its selective antispasmodic action on the digestive tract controls spasticity without the undesirable side effects of atropine or belladonna. Thus, symptomatic relief of many common disturbances of the stomach or intestines can be achieved with discrimination and safety. Mesopin is indicated for the relief of gastrointestinal spasticity, such as pylorospasm, cardiospasm, spastic colon, and biliary spasm.

### MESOPIN

(brand of homatropine methyl bromide)

#### SELECTIVE GASTROINTESTINAL ANTISPASMODIC

SUPPLY: Elixir in 16 ounce bottles; tablets in bottles of 100.

MESOPIN (homatropine methyl bromide)—2.5 mg. per teaspoonful of elixir or per tablet. Also supplied: MESOPIN-PB\*—2.5 mg. Mesopin and 15 mg. (1/4 gr.) phenobarbital per teaspoonful of elixir or per tablet.

Detailed literature and samples on request.

# Endo<sup>®</sup>

\*PB abbreviated designation  
for phenobarbital.

ENDO PRODUCTS INC., RICHMOND HILL 18, NEW YORK

## WASHINGTON LETTER

back the VA hospital construction program.

Agriculture Department is making a mass survey of its 11,000 employees in the Washington area to detect diabetes; active cases are referred to private physicians for treatment.

U.S. Public Health Service has scheduled examinations for medical officers for October 9 through 11. Salaries are \$5,686 and \$6,546.

Five Public Health Service officers have been assigned to help carry out a campaign against malaria and other acute diseases in Indo-China.

Dr. Leonard A. Scheele, U.S. Surgeon General, made a direct appeal to Congress for a higher ceiling on United States appropriations to World Health Organization. He said

WHO—Russia and satellite countries have dropped out—offers a means of carrying on the cold war by improving the health and productivity of noncommunist countries. Just returned from the WHO assembly meeting in Geneva, Dr. Scheele buttressed his argument with facts about the world's health and the WHO record of accomplishment.

Agreement has finally been reached by Food and Drug Administration and retail druggists on a new prescription code.

Most tuberculosis cases among VA staffs do not originate in personnel serving in TB hospitals but in other hospitals.

Start on the new Armed Forces Pathology Institute has been authorized.

**LAVORIS**  
MOUTHWASH AND GARGLE

### Appreciated in the sick room

Tangy  
Cinnamon-clove  
Flavor



ACTIVE INGREDIENTS  
Zinc Chloride - Menthol  
Formaldehyde - Saccharine  
Oil Cinnamon - Oil Cloves  
Alcohol 5%



The thorough cleansing action of Lavoris and its pleasant, refreshing taste are most welcome to the patient.

THE LAVORIS COMPANY MINNEAPOLIS 1, MINN.



TIME-TESTED  
MEDICAMENT



for } **CONSTIPATION  
HYPERACIDITY**

For more than 75 years, Phillips' Milk of Magnesia has been generally accepted by the medical profession as a standard therapeutic agent for constipation and gastric hyperacidity.

*As a laxative*—Phillips' mild, yet thorough action is ideal for both adults and children.

*As an antacid*—Phillips' affords fast, effective relief. Contains no carbonates, hence produces no discomforting flatulence.

**DOSAGE:** { Laxative: 2 to 4 tablespoonfuls  
Antacid: 1 to 4 teaspoonfuls, or  
1 to 4 tablets

**PHILLIPS' MILK OF MAGNESIA**

Prepared only by THE CHAS. H. PHILLIPS CO. DIVISION, 170 Varick Street, New York 13, N. Y.  
of Sterling Drug Inc.

# Current Books & Pamphlets

*This catalogue is compiled from all available sources, American and foreign, to insure a complete listing of the month's releases.*

## Medicine

AKUTE INFektionsKRANKHEITEN UND HOCH-DRUCK: UNTERSUCHUNGEN UND BETRACHTUNGEN ZUM PROBLEM DER POSTINFECTIOSEN HYPERTONIE by Otto H. Arnold. 130 pp. Georg Thieme, Stuttgart. 9.60 M.

A GUIDE TO GENERAL MEDICAL PRACTICE by Martin G. Vorhaus. 244 pp. Macmillan Co., New York City. \$3.50

THE RHEUMATIC DISEASES by G. D. Kersley. 3d ed. 143 pp. William Heinemann Medical Books, London. 15s.

METHODS IN MEDICAL RESEARCH: VOLUME II, edited by Julius H. Comroe, Jr. 361 pp., ill. Year Book Publishers, Chicago. \$6.50

## Pathology

THE 1949 YEAR BOOK OF PATHOLOGY AND CLINICAL PATHOLOGY edited by Howard P. Karsner and Arthur Hawley Sanford. 543 pp., ill. Year Book Publishers, Chicago. \$4.75

## Obstetrics & Gynecology

GYNECOLOGY (THE TEACHINGS OF JOHN I. BREWER) by John I. Brewer. 448 pp., ill. Thomas Nelson & Sons, New York City. \$7.50

A HANDBOOK OF OBSTETRICS AND DIAGNOSTIC GYNECOLOGY by Leo Doyle. 240 pp., ill. University Medical Publishers, Palo Alto, California. \$2

OPERATIVE OBSTETRICS: A GUIDE TO THE DIFFICULTIES AND COMPLICATIONS OF OBSTETRIC PRACTICE by John M. Munro Kerr and J. Chassar Moir. 5th ed. 948 pp., ill. Williams & Wilkins Co., Baltimore. \$12

PRIMARY CARCINOMA OF THE VAGINA by Robert G. Livingstone. 78 pp. Charles C. Thomas, Springfield, Ill. \$2

## Surgery

GARRÉ-STICH-BAUER LEHRBUCH DER CHIRURGIE revised by Rudolf Stich and Karl Henrich Bauer. 14th and 15th ed. 860 pp., ill. Springer, Berlin. 60 M.

## Psychiatry

THE DIAGNOSIS OF HYSTERIA by D. W. Abse. 112 pp. John Wright & Sons, Bristol, England. 8s. 6d.

PSYCHIATRIE: EIN LEHRBUCH FÜR STUDIERENDE UND ÄRZTE by Kurt Kolle. 444 pp. Urban & Schwarzenberg, Berlin. 18 M.

## Radiology

RADIOGRAPHIC ATLAS OF SKELETAL DEVELOPMENT OF THE HAND AND WRIST by William W. Greulich and S. Idell Pyle. 190 pp., ill. Stanford University Press, Stanford, California. \$10

ULTRAVIOLETTE STRAHLEN by A. E. H. Meyer and E. O. Seitz. 2d ed. 390 pp., ill. Walter de Gruyter & Co., Berlin. 34 M.


CLINICAL RADIATION THERAPY edited by Ernst A. Pohle. 2d ed. 902 pp., ill. Lea & Febiger, Philadelphia. \$15

## Orthopedics

OFFICE ORTHOPEDICS by Lewis Cozen. 232 pp., ill. Lea & Febiger, Philadelphia. \$5

KINESIOLOGY: THE MECHANICAL AND ANATOMIC FUNDAMENTALS OF HUMAN MOTION by Katherine F. Wells. 478 pp., ill. W. B. Saunders Co., Philadelphia. \$4.75

THE 1949 YEARBOOK OF ORTHOPEDICS AND TRAUMATIC SURGERY edited by Edward L. Compere. 464 pp., ill. Year Book Publishers, Chicago. \$5



## Your Patient Has 18 to 20 Square Feet of Surface Skin!

The average human body has a surface skin area of 18 to 20 square feet—and every inch is at all times susceptible to one skin disorder or another.

Fortunately, a dermatologic cream exists which is highly effective in alleviating many of these conditions.

## Tarbonis

THE ORIGINAL CLEAN WHITE COAL TAR CREAM

*All the Therapeutic Advantages of Crude Coal Tar with Irritating Residues Removed*

Of 51 difficult dermatologic cases recently treated with TARBONIS in a 5-week to 5-month period, 54.9% cleared or showed marked improvement.\* 25.5% showed good response. TARBONIS brought satisfactory results in 80.4% of the patients! 41 cases involved conditions of 2 to 10 years duration, not yielding to other therapy!

	CASES	CLEARED OR MARKED IMPROVEMENT	MODERATE IMPROVEMENT	SLIGHT OR NO IMPROVEMENT
CHRONIC RECURRENT CONTACT DERMATITIS	11	9	1	1
PSORIASIS	11	2	4	5
NEURODERMATITIS	5	3	2	—
ATOPIC ECZEMA	8	6	1	1
SEBORRHEIC DERMATITIS	6	5	1	—
VARICOSE ECZEMA	4	1	1	2
ALLERGIC DERMATITIS	3	—	2	1
LICHEN PLANUS	3	2	1	—
<b>TOTAL</b>	<b>51</b>	<b>28</b>	<b>13</b>	<b>10</b>
<b>%</b>		<b>54.9</b>	<b>25.5</b>	<b>19.6</b>

For prescriptions—all pharmacies stock 2¼-oz. & 8-oz. jars; for dispensing purposes, 1-lb. & 6-lb. jars available thru your surgical supply dealer.

\*Lowenfish, F.P., N.Y. State J. Med., 50:922 (Apr. 1) 1950.

THE TARBONIS COMPANY Dept. MM  
4300 Euclid Ave., Cleveland 3, Ohio  
Please send literature and clinical sample of TARBONIS

NAME \_\_\_\_\_ M.D.  
ADDRESS \_\_\_\_\_  
CITY \_\_\_\_\_ ZONE \_\_\_\_\_ STATE \_\_\_\_\_



**further  
improvement  
in postpartum  
breast care  
technic**

New nylon Plastishields®  
may be sterilized by boiling  
or autoclaving!


- Save nurses' time . .
- Encourage Breast feeding...
- Reduce incidence of sore,  
cracked and fissured nipples  
... with Plastishields!

Pat. No. 2495307

**Plastishield, inc.**

Minneapolis 3, Minnesota

**R<sub>x</sub>** FOR YOUR  
OFFICE



Gleaming White Porcelain  
and Stainless Steel

**RENWAL<sup>®</sup>  
STERILIZERS**

Automatic  
Shut-Off Valve

Easy to Keep Clean  
Won't Rust

No. 12—11 inch \$29.50  
No. 18—7 1/2 inch 13.50  
No. 5—5 inch 7.95

At All Surgical Dealers

**R<sub>x</sub>** FOR YOUR  
DIABETIC  
PATIENTS



\$7.95

**RENWAL No. 5<sup>®</sup>  
DIABETIC  
STERILIZER**

Specially Designed for  
Personal and Private Service

For Syringes, Needles  
and Urine Tests

At All Surgical Dealers

**AMERICAN SUNDRIES CO., INC., B'KLYN, N. Y.**

Patented

**Dermatology**

THE 1949 YEARBOOK OF DERMATOLOGY AND  
SYPHILOLOGY edited by Marion B. Sulz-  
berger and Rudolf L. Baer. 514 pp.,  
ill. Year Book Publishers, Chicago. \$5

**Pharmacology**

DRUG PLANTS OF AFRICA by Thomas S.  
Githens. 125 pp. University of Penn-  
sylvania Press, Philadelphia. \$2.25

ANGEWANDTE PHARMAKOLOGIE FÜR ÄRZTE  
UND STUDIERENDE DER MEDIZIN by Hesse  
3d ed. 445 pp., ill. Urban & Schwarz-  
enberg, Berlin. 23 M.

A CONCISE APPLIED PHARMACOLOGY AND  
THERAPEUTICS by F. G. Hobart and G.  
Melton. 3d ed. 300 pp. Leonard Hill.  
London. 21s.

**Nursing**

METHODS OF TEACHING IN SCHOOLS OF  
NURSING by Alice B. Brethorst 362 pp.,  
ill. W. B. Saunders Co., Philadelphia.

\$4

NURSING HISTORY IN BRIEF by Minnie  
Goodnow. 3d ed. 274 pp., ill. W. B.  
Saunders Co., Philadelphia. \$3

MEDICAL NURSING by Edgar Hull and  
Cecilia M. Perrodin. 4th ed. 826 pp.,  
ill. F. A. Davis Co., Philadelphia. \$4

THE SOCIOLOGY OF THE PATIENT: A TEXT-  
BOOK FOR NURSES by Earl Lomon Koo. 264  
pp. McGraw-Hill Book Co., New  
York City. \$3

A PRACTICAL HANDBOOK OF PSYCHIATRY  
FOR STUDENTS AND NURSES by L. Minski.  
2d ed. 136 pp. William Heinemann  
Medical Books, London. 6s.

MEDICINE FOR NURSES by William Gor-  
don Sears. 5th ed. 472 pp., ill. Wil-  
liams & Wilkins Co., Baltimore. \$3

**THE BEE CELL**

Reg. U. S. Pat. Off.

**A PESSARY OF  
PROVEN EXCELLENCE**

Made from finest, soft, natural  
rubber. Easily removed and  
replaced by wearer. Worn with ease and comfort.  
Secures support by suction of six concave surfaces.

Sold directly to physicians or on the prescription  
of a physician—not through dealers.

Information, descriptive literature and reprints\*  
mailed to physicians on request.

\*The Bee Cell pessary. West. J. Surg. 57: 481-482,  
1949

**THE BEE CELL CO.**

P. O. Box 212 Dept. A Buffalo 5, N. Y.




NOW!



CHILDREN'S SIZE

*Genuine*   
**BAYER**  
ASPIRIN

This special Children's Size Bayer Aspirin makes it easy for mothers to follow your prescribed dosage.



Write for a package  
The **BAYER COMPANY DIVISION**  
OF STERLING DRUG INC.  
170 Varick Street • New York 13, N. Y.





## HAY FEVER?

**Perfumed Cosmetics Can Induce Symptoms**

Many physicians routinely prescribe **AR-EX Unscented Cosmetics**. Eliminate a whole field of respiratory sensitizers. Fashion-right shades. Pleasant to use. Beautifully packaged.

Send for Free Formulary  
**AR-EX COSMETICS, INC.**  
1036 W. Van Buren, Chicago 7

**AR-EX UNSCENTED COSMETICS**



**Write for Sample**

The Alkalol Company, Taunton 10, Mass.

**WHEN TREATMENT IS INDICATED —**

**RECOMMEND** → **THUM** TRADE MARK

To Discourage **NAIL-BITING**

**PAINT ON FINGERTIPS**

USE THUM IN STUBBORN THUMB-SUCKING CASES TOO

**50c and \$1.00 ORDER FROM YOUR SUPPLY HOUSE OR PHARMACIST**



## PATIENTS

### ... I Have Met

The editors will pay \$1 for each story published. No contributions will be returned. Send your experiences to the *Patients I Have Met* Editor, MODERN MEDICINE, 84 South Tenth St., Minneapolis 3, Minn.

#### Inevitable Progression

The doctor had advised the man to slow down in his living. First the patient cut out tea, then coffee, smoking, and drinking. Next he cut out women, and now he's cutting out paper dolls.  
—I.H.O.

"What I want," said the patient who had just undergone an operation for ischiorectal fistula, "is a pretention enema."—P.W.V.



"Robber! Thief! Bandit! Go get yourself a pair of glasses."

#### Willing to Help

A friend of mine decided to send his fourteen-year-old son to a prep school but felt that he first should have a talk with him on the subject of sex.

"Now that you are going away to school, Willy," he said, "I feel that I should discuss with you what I consider a very important matter. I am at a loss, however, as just how to start, not knowing how much you know about sex."

"That's o.k., dad," responded the son. "What do you want to know about it?"

—A.H.M.

# AUREOMYCIN

CRYSTALLINE

## *in Infections of the Puerperium*

During the past year, obstetricians have become increasingly impressed with the ability of aureomycin to prevent or arrest infections of the puerperium. Where infection is feared, or has appeared, this broadly effective antibiotic is highly useful. Drug fastness and allergy are very rare following aureomycin. It is believed that this new crystalline form of aureomycin obviates nearly all side reactions.

Aureo-  
mycin



Aureomycin has also been found effective for the control of the following infections:

Acute amebiasis, bacterial infections associated with virus influenza, bacterial and virus-like infections of the eye, bacteroides septicemia, boutonneuse fever, brucellosis, chancroid, Friedländer infections (*Klebsiella pneumonia*), gonorrhea (resistant), Gram-negative infections (including those caused by some of the coli-aerogenes group), Gram-positive infections (including those caused by streptococci, staphylococci, and pneumococci), granuloma inguinale, *H. influenzae* infections, lymphogranuloma venereum, peritonitis, pertussis infections (acute and sub-acute), primary atypical pneumonia, psittacosis (parrot fever), Q fever, rickettsialpox, Rocky Mountain spotted fever, sinusitis, subacute bacterial endocarditis resistant to penicillin, surgical infections, tick-bite fever (African), tularemia, typhus and the common infections of the uterus and adnexa.

### **Capsules:**

Bottles of 25, 50 mg. each capsule.  
Bottles of 16, 250 mg. each capsule.

### **Ophthalmic:**

Vials of 25 mg. with dropper;  
solution prepared by adding  
5 cc. of distilled water.



# ARTHRITIS



One Gelucap Weapon For 2-Way Therapy

Many cases of arthritis are accompanied by fibrositis. Steinsberg\* showed 143 out of 145 cases of primary fibrositis aided by high potency vitamin E. EDREX offers both vitamin D and vitamin E. (\*Am. Inst. Med., 19:136-139) Send for free literature.

## EDREX

VITAMINE  
PLUS  
VITAMIN D

WILCO LABORATORIES  
800 N. Clark St., Chicago 10, Ill.




## Mechanical Aid In Relief Of Allergic Suffering

As an adjunct to your treatment of inhalant allergic symptoms, recommend the use of the Weaver Nasal Filter. Filters out many possible allergens and irritants from the air breathed, individually fitted to patient's nostrils by trained technicians. Sanitary silver cups with replaceable filter mats are comfortably worn and are scarcely visible. For literature and address of nearest fitting agency, write

NASAL FILTER CO., Colum<sup>15</sup>, Chi

Dept. B1

## This GOMCO Pump for SAFE Administration of Anti-Biotics...



• The Gomco No. 788-0 AEROSOL PENICILLIN PUMP provides ACCURATELY REGULATED compressed air for atomization of solutions as desired... for safe, positive administration of penicillin, streptomycin, other anti-biotics. A sturdy, precision-built pump for many years of service. ASK YOUR DEALER OR WRITE US, requesting complete GOMCO catalog.

Gomco Surgical Manufacturing Corp.  
844M E. Ferry St. Buffalo 11, N.Y.

## For the Record

My new secretary, a very conscientious girl, was puzzled by an entry in my notes on an emergency case: "Shot in the lumbar region." After a moment's study she brightened, hitched her chair up to the typewriter and typed into the record, "Shot in the woods."—F.M.S.

"I just love orchids," said the night nurse. "Just a case of orchiditis, I suppose."—B.S.

## Time for a Dry Spell

I had given a new mother with vaginitis a prescription for an astringent douche. Two weeks later she called to report that everything was all right. In that case, I said, she could stop the douches.

"Thank you, doctor," she said. "Between the douches, the dishes, and the diapers, I've had my hands in water all the time."—F.M.M.

"Doctor," complained the mother of a month-old baby boy, "he gets awfully sore under his genitals."—M.L.R.

## Waiting Room Amnesia

It had been a long weary session of evening office hours. Finally I got around to calling a young lady who had been waiting patiently for some time. As she sat down in the chair by my desk, she sighed, "Doctor, I been waiting so long I forgot what I came for."—P.C.J.



## LITTLE TOIDEY for Training the Baby

Little Toidey, \$5.50, and Toddler's Toidey (base and pan) \$3.95, are most convenient in the office of physicians who have patients with babies. 20% COURTESY DISCOUNT to doctors on all orders mailed directly to us. Write for complete list and free book "Training the Baby."

## THE TOIDEY COMPANY

Gertrude A. Muller, Inc.  
FORT WAYNE • INDIANA

#### NURSING ARTS,

Mildred L. Montag, M.A., R.N.,  
Margaret Filson, M.A., R.N.,  
Saunders, 1948: p. 237—

"Back care cannot be over-emphasized."

p. 377—

"The practice of rubbing the skin, particularly the back, with alcohol to prevent pressure sores is not altogether logical. Alcohol is drying and a dry skin is more susceptible to cracking and irritation than is a somewhat oily skin surface . . . Therefore, some kind of lubricant, such as a lotion, seems to be indicated in the care of the back."



## HOW IMPORTANT IS PROTECTION of *the Bed Patient's Skin?*

DERMASSAGE is a soothing emollient cream, superior to alcohol for body massage, for use after bathing, and in measures for the prevention of bedsores. Water soluble lanolin (safe for use of allergic persons) and olive oil form the base for absorption. Contains also natural menthol U. S. P.; oxyquinoline sulphate, a non-toxic antiseptic and bacteriostatic having deodorant properties; and carbamide, whose solvent action on proteins aids removal of debris and dead tissue.

The singular function of DERMASSAGE is the protection of the skin of the hospitalized patient. Your supplier has DERMASSAGE.

# EDISON'S dermassage

USED IN OVER  
4,000 HOSPITALS

Advertised in  
American Medical  
Association  
Publications

You are invited to test DERMASSAGE firsthand  
EDISON CHEMICAL COMPANY  
30 W. Washington St., Chicago 2

Please send me, without charge, your PROFESSIONAL  
SAMPLE of Edison's DERMASSAGE.

Name \_\_\_\_\_

Address \_\_\_\_\_

MM 8-50



**RECESSED SPECIALIST'S OUTFIT**  
See it at your dealer, or write

**PROMETHEUS**

**ELECTRIC CORPORATION**  
401 W. 13th St. New York 14

**Gebauer's**  
**S.S. SPRAY**



## New... Instant Detergent Lather

• This unique development provides an instantaneous, rich, creamy lather for preparing the field of operation prior to use of antiseptic in minor surgery...removal of ointment dressings...rapid removal of oil and grease in industrial injuries...removing oily base in syringes. Efficient, effective, economical. At your surgical supply house, or write for literature.

**THE GEBAUER CHEMICAL COMPANY**  
9410 St. Catherine Ave. • Cleveland, Ohio

## INDEX TO ADVERTISERS

*The publishers are not responsible  
for any errors or omissions.*


Abbott Laboratories.....	83
Alkaloi Co., The.....	110
American Ferment Company, Inc.....	15
American Sundries Co., Inc.....	168
Ar-Ex Cosmetics, Inc.....	110
Barnes, A. C. Company.....	13
Bauer & Black.....	38
Bausch & Lomb Optical Co.....	93
Bayer Aspirin.....	169
Ree Cell Co., The.....	168
Belmont Laboratories.....	79
Borden Company, The.....	60-61
Burroughs Wellcome & Co., Inc., (U.S.A.).....	16
Central Pharmacal Company, The.....	101
Cereal Institute, Inc.....	86-87
Chatham Pharmaceuticals, Inc.....	89
Chilcott Laboratories.....	5
Ciba Pharmaceutical Products, Inc.....	37, 4th Cover
Colwell Publishing Co.....	102
Eastman Kodak Company.....	32-33
Edison Chemical Company.....	113
Endo Products Inc.....	103
Gebauer Chemical Company, The.....	114
Geriatrics.....	116
Gomco Surgical Manufacturing Corp.....	112
Hoffmann-La Roche, Inc.....	10
Johnson & Johnson.....	22-23
Knox Gelatine.....	25
Kremers-Urban Company.....	39
Lavoris Company, The.....	104
Lederle Laboratories, Inc.....	111
McNeill Laboratories, Inc.....	3
Merrell, The Wm. S. Company.....	2nd Cover, 6-7, 75, 91
Nasal Filter Co.....	112
National Drug Company, The.....	31
Num Specialties.....	110
Pfizer, Chas. & Co., Inc.....	40-41
Phillips Co., The Chas. H.....	105
Pioneer Rubber Co., The.....	36
Plastisfield, Inc.....	108
Professional Printing Company, Inc.....	26
Prometheus Electric Corp.....	114
Ritter Co., Inc.....	9
Robins, A. H., Company, Inc.....	4, 96-97
Schenley Laboratories, Inc.....	84
Schleffelin & Co.....	90
Searle & Co., G. D.....	80-81
Sherman Laboratories.....	77
Shield Laboratories.....	28-29
Smith, Kline & French Laboratories.....	17, 27, 99
Smith, Martin H., Company.....	92
Stuart Company, The.....	19, 20
Talby-Nason Company.....	21
Tarbons Company, The.....	107
Tolidey Company, The.....	112
Ulmer Laboratories.....	115
Varick Pharmacal Co., Inc.....	34
White Laboratories, Inc.....	35, 95
Wilco Laboratories, The.....	112
Winthrop-Stearns Inc.....	3rd Cover
Wyeth Incorporated.....	42

FOR THE DOCTOR'S OFFICE PRACTICE

# BILIPOSOL

TRADE MARK

REG. U. S.  
PAT. OFF.



A  
Special  
LIPOID SOLUBLE  
BASIC BISMUTH  
in clear  
homogeneous  
oil solution  
FOR  
SUCCESSFUL  
TREATMENT  
OF ALL STAGES  
OF  
**SYPHILIS**

## ★ Painless Intramuscular Injection

## ★ RAPID SPIROCHAETICIDAL ACTION

THE TREPONEMA USUALLY DISAPPEAR FROM THE LESIONS  
IN 24 HOURS FOLLOWING THE FIRST INJECTION

## ★ EARLY NEGATIVE WASSERMANN

## ★ IMMEDIATE ABSORPTION WITH SLOW ELIMINATION

Renal elimination of bismuth commences rapidly and three hours after injection the metal is demonstrable in the urine. It continues after cessation of the treatment up to 1½ to two months.

## BILIPOSOL HAS STOOD THE TEST OF TIME

BILIPOSOL, in Boxes of 12, 50 and 100 ampoules, each containing 8 cty. bismuth,  
IS OBTAINABLE FROM LEADING PHYSICIANS' SUPPLY  
HOUSES, RETAIL AND WHOLESALE DRUGGISTS OR FROM

*Sole General Distributor for United States and Canada*

## ULMER LABORATORIES

Minneapolis Minnesota

ULMER PHARMACAL COMPANY, Northwest Distributors

412 South Sixth Street, Minneapolis, Minnesota

Literature to physicians on request

DEVOTED TO RESEARCH AND CLINICAL  
DISEASES AND PROCESSES OF THE AGED

# Geriatrics

Official Journal of the  
AMERICAN GERIATRICS SOCIETY

Join the thousands of your colleagues who read GERIATRICS for the most significant developments in research and clinical study of the diseases of the aged and aging. GERIATRICS keeps you abreast of progress in diagnosis and treatment of cardiovascular, neurological, arteriosclerotic, arthritic, proctologic, prostatic, nutritive and other disorders.

Subscribe NOW! — Save \$5 on BARGAIN  
INTRODUCTORY OFFER — 3 YEARS, \$10!

GERIATRICS, 84 South 10th St., Minneapolis 3, Minn.

I'd like to have GERIATRICS help me keep abreast of latest developments in geriatrics. I'll take your bargain subscription deal.

Name.....Address.....

City.....Zone.....State.....

Check Enclosed..... Please Bill Me.....

GERIATRICS is chosen by the most eminent authors to present their important papers on geriatrics.



when edematous mucous membranes  
cause nasal obstruction...

**NEO-SYNEPHRINE**  
**HYDROCHLORIDE**  
BRAND OF PHENYLEPHRINE HYDROCHLORIDE

*Winthrop Stearns* INC.  
NEW YORK 12, N. Y. WINDSOR, ONT.

**SUPPLIED AS**

¼ % and 1 % strengths  
in isotonic saline solution—  
1 oz. bottles.

¼ % in isotonic solution  
of three chlorides (Ringer's) with  
aromatics—1 oz. bottles.

½ % water soluble jelly—  
½ oz. tubes.

in hay fever—the vasoconstrictive effect  
of Neo-Synephrine hydrochloride  
¼ % to 1 % on the engorged mucous  
membrane is rapid and prolonged—but  
with virtually no discomfort.

The patient quickly experiences  
welcome relief of the "stuffy" feeling,  
breathing is eased,  
drainage is promoted,  
and general irritation is lessened.  
Neo-Synephrine hydrochloride is a  
valuable aid in circumventing  
sleeplessness which may be associated  
with hay fever—for its protracted  
shrinking effect on the nasal  
membranes overcomes the annoyance  
of "mouth breathing."

Neo-Synephrine hydrochloride  
exerts an undiminished effect after  
repeated use; it usually does not  
produce compensatory congestion,  
and has virtually no central  
stimulating action.


Neo-Synephrine, trademark reg. U. S. & Canada

# New Method for Relief of Allergic Nasal Symptoms

This convenient plastic Nebulizer distributes a mist of minute droplets of PYRIBENZAMINE hydrochloride Nasal Solution throughout the nasal passages.

Relief usually is immediate—complete—prolonged. Side reactions rarely occur except for occasional transient stinging. It is convenient to carry in purse or pocket and may be used at any time in any place.

The Nebulizer provides several hundred applications of PYRIBENZAMINE hydrochloride 0.5% in an isotonic, buffered solution. One application in each nostril usually is a therapeutic dose and may be repeated as required.



## Pyribenzamine Nebulizer

**Ciba** PHARMACEUTICAL PRODUCTS, INC.  
SUMMIT, N. J.

PYRIBENZAMINE® (brand of tripeleminamine) 2/1954M

MODERN MEDICINE  
84 S. 10 St. Minneapolis 3, Minn.

POSTMASTER: If undeliverable FOR ANY REASON  
notify sender, stating reason on form 3547.

Sec. 34.66, P. L. & R.  
U. S. POSTAGE  
**PAID**  
PERMIT No. 29  
Long Prairie, Minn.